

SWALLOWING THE HURT: EXPLORING THE LINKS BETWEEN FAMILY VIOLENCE, ANOREXIA, AND BULIMIA

Prepared by

Shelley Moore, Ph.D.

With the assistance of

Kelly D'Aoust, B.A.

Yasmin Jiwani, Ph.D.

Donna Robertson, B.A.

Christina Savage, B.A.

for

The FREDA Centre for Research on Violence against Women and Children
Vancouver, BC

Submitted to

Family Violence Prevention Unit
Health Issues Division
Health Canada

October 2001

TABLE OF CONTENTS

| | |
|--|----|
| EXECUTIVE SUMMARY | 1 |
| PURPOSE AND BACKGROUND | 3 |
| PRODUCING THE REPORT: THE ADVISORY PANEL AND SERVICE PROVIDERS | 7 |
| <i>METHOD</i> | 7 |
| <i>APPROACH</i> | 8 |
| THE NATURE OF VIOLENCE AGAINST WOMEN AND GIRLS AND OF EATING DISORDERS IN CANADA | 11 |
| <i>PREVALENCE</i> | 11 |
| Violence Against Women and Girls | 11 |
| Eating Disorders | 12 |
| Overall prevalence of eating disorders | 12 |
| Prevalence rates for bulimia | 12 |
| Prevalence rates for anorexia..... | 13 |
| Breaking prevalence rates down..... | 13 |
| Diagnostic criteria | 14 |
| <i>IMPACT</i> | 15 |
| <i>GIRLS AND WOMEN DYING</i> | 15 |
| LINKS BETWEEN VIOLENCE AGAINST WOMEN AND GIRLS, ANOREXIA, AND BULIMIA | 17 |
| <i>ABUSE OF CHILDREN AND THE DEVELOPMENT OF EATING DISORDERS</i> | 18 |
| Childhood Sexual Abuse | 18 |
| Childhood Physical Abuse..... | 19 |
| Family Interactions during Childhood..... | 19 |
| <i>YOUTH AND ADOLESCENT EXPERIENCES OF VIOLENCE, ANOREXIA, AND BULIMIA</i> | 20 |
| <i>VIOLENCE AND EATING DISORDERS IN ADULTHOOD AND AGING</i> | 23 |
| Battering | 23 |
| Sexual Trauma in Adulthood..... | 24 |
| Aging | 25 |
| <i>THE ROLE OF DISCLOSURE</i> | 26 |
| <i>AGE</i> | 27 |
| UNDERSTANDING THE LINKS | 29 |
| <i>CAUSALITY: DO EXPERIENCES OF VIOLENCE LEAD TO EATING DISORDERS?</i> | 29 |
| <i>COPING WITH VIOLENCE</i> | 29 |
| <i>CONTROL AND POWER</i> | 31 |
| <i>SELF-ESTEEM</i> | 32 |
| <i>THE SOCIOCULTURAL CONTEXT</i> | 33 |
| ISSUES IN TREATING VIOLENCE AND EATING DISORDERS | 35 |
| <i>INTEGRATION OF TREATMENT</i> | 35 |
| <i>SELF-ESTEEM</i> | 36 |
| <i>INDIVIDUALIZING TREATMENT</i> | 36 |
| <i>BARRIERS OF MARGINALITY</i> | 37 |
| <i>GEOGRAPHIC ISOLATION</i> | 39 |
| <i>SAFETY</i> | 40 |
| <i>DEVELOPMENTAL TREATMENT ISSUES</i> | 40 |
| <i>LACK OF FUNDING AND SERVICES</i> | 41 |
| KEY COMPONENTS IDENTIFIED FOR THE TREATMENT OF VIOLENCE AND EATING DISORDERS | 43 |
| <i>PRESENTATION</i> | 44 |
| 1. Accessibility | 44 |
| 2. Assessment | 44 |
| <i>IMMEDIATE NEEDS</i> | 45 |
| 3. Safety | 45 |
| 4. Medical Stability..... | 45 |
| <i>THERAPEUTIC SUPPORT</i> | 46 |

| | |
|--|----|
| 5. Validation | 46 |
| 6. Self-worth | 46 |
| 7. Re-framing..... | 46 |
| 8. Coping Skills | 46 |
| 9. Control and Empowerment..... | 46 |
| 10. Relationships and Networks | 47 |
| 11. Education and Awareness | 47 |
| <i>CONTINUITY</i> | 47 |
| 12. Follow-up | 47 |
| CONCLUSION..... | 49 |
| WORKS CITED | 51 |
| APPENDIX A: THE ADVISORY PANEL..... | 67 |
| APPENDIX B: A LIST OF PROGRAMMES | 69 |
| <i>BRITISH COLUMBIA</i> | 69 |
| <i>YUKON</i> | 73 |
| <i>NORTHWEST TERRITORIES</i> | 73 |
| <i>ALBERTA</i> | 74 |
| <i>SASKATCHEWAN</i> | 75 |
| <i>MANITOBA</i> | 76 |
| <i>ONTARIO</i> | 77 |
| <i>QUÉBEC</i> | 81 |
| <i>NEW BRUNSWICK</i> | 82 |
| <i>PRINCE EDWARD ISLAND</i> | 83 |
| <i>NOVA SCOTIA</i> | 84 |
| <i>NEWFOUNDLAND</i> | 86 |
| APPENDIX C: INTERVIEW QUESTIONS..... | 87 |

LIST OF FIGURES

| | |
|--|----|
| FIGURE 1: RESPONSES BY CANADIAN SERVICE PROVIDERS TO A POSSIBLE LINK BETWEEN FAMILY VIOLENCE AND EATING DISORDERS | 17 |
|--|----|

EXECUTIVE SUMMARY

This report examines the links between eating disorders and violence against women and girls. It is based on information gathered from published literature as well as consultations with community workers, health practitioners, and mental health professionals. "Swallowing the Hurt" has been designed for use by front-line workers, health-care and social service professionals, educators, and researchers who offer services directly for or who may interact with women and girls experiencing eating disorders or violence.

Over the last decade increasing attention has been turned to anorexia and bulimia as possible outcomes of abuse. Both the literature and most of the service providers contacted suggest that sexual, physical, verbal, and emotional abuses are contributing factors in a complex and multideterminant model of anorexia and bulimia. The research suggests that the connection between violence and eating disorders may be more pronounced when: food has been used as a weapon of abuse; a woman has been abused by more than one person; disclosure has been punished or disbelieved; multiple forms of abuse have been experienced; or the woman or girl feels a greater sense of powerlessness or shame.

It is estimated that at least 1 in 8 Canadian girls will be seriously sexually abused and 1 in 5 will be physically abused before the age of 18 years (Duffy, 1998). At least 10% to 25% of women will be battered by their husbands or common-law spouses (Benson, 1995; Statistics Canada, 1993). Abuse rates are 8-times higher for Aboriginal women (Duffy, 1998) and 4 times higher for women with disabilities (Razack, 1994). The psychological and health impacts of violence against women and girls can be serious and enduring. Eating disorders represent particularly life-threatening consequences for survivors of abuse. Across a 20-year period, 1 in 5 people with anorexia and at least 5% of people with bulimia will die (Woodside, 1995). Girls and women constitute 90% of those fatalities (Gagnon, 1996). In addition, 56% of those with anorexia experience severe medical illnesses (Herzog, Deter, Fiehn, & Petzold, 1997). Both violence and eating disorders affect predominantly girls and women and both appear to be rising in prevalence (American Psychiatric Association, 1994; Hsu & Zimmer, 1988).

Of those service providers who are aware of the links between eating disorders and abuse, not all are able to offer an integrated approach to treatment. Some refer the woman/girl to other agencies or out of province; some only have resources to focus on a single issue. Both the service providers and the advisory panel for this project emphasized a greater need for "cross-training" between the fields of disordered eating and violence. From the research and the consultations, 12 key components have been identified as optimal to an integrated approach: accessibility; assessment; safety; medical stability; validation; self-worth; re-framing; coping skills; control and empowerment; relationships and networks; education and awareness; and follow-up. A primary concern expressed by service providers, however, is the effect of cutbacks on funding and facilities. Programmes are suffering a loss of staff and services, a lack of follow-up care, increasing wait lists, and closures. In many cases, the need for funding has become urgent. Without the resources to ensure accessibility to services, appropriate medical care, sufficient staffing, shelters and

transition houses, and counselling for underlying abuse issues, integrated treatment approaches to family violence and eating disorders are likely to become increasingly scarce.

It should be emphasized that although this report explores the links specifically between violence and the development of disordered eating, the vast majority of participating service providers cautioned that abuse should not be considered as the only precursor to eating disorders.

PURPOSE AND BACKGROUND

Past research has demonstrated that women with more conflictive family backgrounds or childhood experiences are more likely to develop eating problems (Kinzl, Traweger, Guenther, & Biebl, 1994), particularly bulimia (Schmidt, Tiller, & Treasure, 1993). There is an increasing amount of research being conducted to explore the role that abuse history might play in the development of eating disorders. Reviews examining this association have been published since 1992 (Wonderlich, Brewerton, Jolic, Dansky, & Abbott, 1997). Overall, the results of the research have been inconsistent (Connors & Morse, 1993; Mullen, Martin, Anderson, Romans, & Herbison, 1993; Reto, Dalenberg, & Coe, 1996) and controversy has emerged among theorists. In 1997, Brown observed that "over the last 5 years, the scientific literature has become polarized, with a heated, and sometimes vitriolic, debate played out between recognized experts, within and across countries" (p. 194). Whereas some researchers espouse a causal relationship between abuse and eating disorders (Everill & Waller, 1995a), others have characterized the link as neither special nor specific (Pope & Hudson, 1992) or as coincidental (Finn, Hartman, Leon, & Lawson, 1986). Most theorists, however, have argued that the relationship is a complex one that should not be over-simplified (Brown, 1997; Welch & Fairburn, 1996; Wonderlich et al., 1997).

Much of the inconsistency in the literature has been attributed to methodological and interpretive difficulties. Widely varying definitions of abuse, and particularly child sexual abuse, have made comparisons among studies and conclusions difficult (Dansky, Brewerton, Kilpatrick, & O'Neil, 1997; Miller, 1996). Whereas some researchers define child sexual abuse as both contact and noncontact experiences (Moyer, DiPietro, Berkowitz, & Stunkard, 1997), others measure only those experiences that include physical contact (Schmidt, Tiller, & Treasure, 1993). Diagnostic criteria of anorexia and bulimia have also been inconsistent across studies (Connors & Morse, 1993; Davis & Yager, 1992; Wiederman, 1996). The criteria for anorexia nervosa and bulimia nervosa within the *Diagnostic and Statistical Manual-IV (DSM-IV)*¹ of the American Psychiatric Association have been revised three times, making comparison across studies over time problematic (Hof & Nicolson, 1996). Moreover, there has been disagreement regarding which symptoms should be required for diagnosis. Amenorrhea (cessation of menstruation) has been contested as neither a necessary nor useful criterion in determining anorexia (Andersen & Holman, 1997; Cachelin & Maher, 1998; Garfinkel et al., 1995), and as one that excludes diagnosis of pre-adolescent girls (Woodside, 1995) and post-menopausal women (Cosford & Arnold, 1992). A number of researchers have critiqued fear of fatness as a necessary criterion for diagnosis, noting in particular that it may be a culturally specific symptom (Hsu & Lee, 1993; Lee, 1995; Lee, Chiu, & Chen, 1989; Thompson, 1992), and a relatively recent addition to definitions of eating disorders (Hof & Nicolson, 1996; Palmer, 1993; Parry-Jones & Parry-Jones, 1995). Others have emphasized eating problems as a

¹ The *Diagnostic and Statistical Manual-IV* is the most recent edition of the text published by the American Psychiatric Association for the purpose of categorizing and clinically describing psychiatric disorders. It is intended for use in research and professional diagnosis.

continuum, much of which may be missed through a strict adherence to diagnostic classifications (Hoek, 1995; Zerbe, 1992). Overall, *DSM-IV* criteria may fail to detect a large number of individuals in significant distress due to disordered eating.

The research on abuse and eating disorders has shifted from an earlier period of case studies to more recent reports of larger studies (Welch & Fairburn, 1996). Nonetheless, current research has continued to rely heavily on clinical (Moyer et al., 1997) and small (Wiederman, 1996) samples. Results have often reflected the type of sample and methodology used. Uncontrolled correlational studies of community samples have generally shown more support for a connection than have controlled examinations of abuse within women with eating disorders or of eating problems among abuse survivors (Connors & Morse, 1993; Wonderlich et al., 1997). Because both abuse rates and eating problems are higher in clinical samples, reviewers have questioned whether significant correlations are more artifactual than meaningful (Connors & Morse, 1993). Moreover, it is not clear whether abuse rates are higher among women with eating disorders than among women with other psychiatric or mental health difficulties (Everill & Waller, 1995b; Herzog, Staley, Carmody, Robbins, & van der Kolk, 1993; Wonderlich et al., 1997), particularly depression, anxiety disorders, somatization, and complex personality disorders (Welch & Fairburn, 1996). But, research has also suggested that eating disorders tend to be more enduring than many other psychiatric disorders, and that repeated, severe abuse may lead to more chronic manifestations of mental health problems (Herzog et al., 1993; Welch & Fairburn, 1996). Herzog et al. (1993) have noted that the co-occurrence of eating disorders and other mental health problems, as well as the comparable rates of abuse among women with other mental health problems, challenges the specificity of the relationship between abuse and troubled eating. Although the uniqueness of the relationship may be in question, however, it cannot be subsequently concluded that past abuse is a meaningless factor in clinical practice.

A second wave of studies with tightened methodology (Wonderlich et al., 1997) has led to greater agreement among researchers. Reviewers have acknowledged that a complete understanding of the relationship between abuse and disordered eating has not yet been attained (Brown, 1997; Everill & Waller, 1995b). As a result, current models emphasize the need for a multidimensional approach in which abuse is not treated as a sole or exclusively causal factor of anorexia and bulimia (Connors & Morse, 1993; Dansky et al., 1997) and eating disorders are recognized as one possible outcome of abuse (Schaaf & McCanne, 1994). The impact of abuse will vary according to the context in which it occurs, the meaning for the individual survivor, and the resilience that may result from optimal family or other supporting relationships (Connors & Morse, 1993; Welch, Doll, & Fairburn, 1997). Overall, Brown (1997) has concluded that researchers do not generally dispute that history of abuse is neither necessary nor sufficient for the development of disordered eating. Studies are increasingly examining the contributory factors that may determine the route from abuse to troubled eating (Everill & Waller, 1995b). This study was a part of that project.

The first purpose of this report was to further explore the existence, strength and nature of a connection between abuse and eating disorders as experienced by service providers across Canada. In particular, we asked service providers the extent to which they witnessed a link and their explanations for the presence or absence of a link. A second purpose was to integrate and to compare the experiential data gathered from Canadian service providers with the clinical and empirical data published in the literature. A third purpose was to examine the connection across the lifespan, with particular attention to the unique developmental challenges of childhood, adulthood, and aging. Using such an inclusive framework, we also found the need for a broader definition of violence. The report therefore includes references to sexual abuse, physical abuse, emotional abuse, battering, sexual harassment, verbal harassment, sexual assault, neglect, family dynamics, systemic violence, and institutional violence. A fourth purpose was to use the feedback received from service providers to determine the need for and to develop an integrated treatment model for violence and disordered eating.

PRODUCING THE REPORT: THE ADVISORY PANEL AND SERVICE PROVIDERS

METHOD

This report is based on a review of published literature and on information gathered through consultations with practitioners. The authors would like to thank the many service providers and the advisory panel for their participation, willingness to share knowledge, and their contributions toward this project.

In producing this report, we contacted 143 service providers across Canada working in the areas of violence and eating disorders. Of these, 123 responded and agreed to participate in the study. Service providers were identified for contact through referrals, Internet searches, community listings, published research in journals and reports, and telephone directories. The service providers who participated offer support through private practice (10%), family or mental health services (15%), public education or outreach programmes (6%), hospitals or health units (25%), treatment and support organizations (32%), community services (6%), general child and youth programmes (4%), and university or college services (2%). We gathered information through informal telephone consultations and discussions with informants in BC (39), NWT/Nunavut (2), Alberta (12), Saskatchewan (16), Manitoba (4), Ontario (34), Québec (9), New Brunswick (4), Prince Edward Island (3), Nova Scotia (15), and Newfoundland (3). The service providers were informally interviewed about possible connections between family violence and eating disorders, about appropriate treatment approaches in addressing these connections, and about their particular services and clientele. Six standard questions were used as a departure point for the interviews (See Appendix C). Participants were encouraged to elaborate on information that they felt was important to convey. Responses were accepted by telephone, fax, or e-mail in order to maximize response rate. The resulting data were analyzed for dominant themes, compared to the published literature, and used to develop an integrated treatment model. In order to protect the anonymity of the service providers, names, geographical locations, and affiliations have been omitted in citing comments and data.

For the final production of this report, a panel of 11 Canadian advisors was assembled to participate in a one-day teleconference held in November 1999 to further explore links between violence against women and girls, anorexia, and bulimia. The suggestions generated during this consultation were integrated into the text of the report. Additionally, comments were elicited individually from treatment providers who were unable to participate in the teleconference. Advisory panel members were selected based both on their recommendation by other service providers as Canadian experts in the fields of disordered eating and violence against women, and on their geographic location.

APPROACH

We took the following approach in undertaking and presenting this research:

- This report is a qualitative analysis that provides both: (a) direct quotes from participants where they contribute insight, precision, or substantiation; and (b) frequencies with which responses were given, to help the reader differentiate the perspective of a single service provider from those of multiple service providers. The inclusion of frequencies within qualitative analysis is common within social science publications (cf. Pope & Vetter, 1992, published in the APA journal, *American Psychologist*, and Jiwani, 1998, prepared for the Department of Justice).
- This document reports the direct responses of service providers to questions regarding the presence and strength of a link between abuse and disordered eating. In addition, it provides a thematic analysis of the experiences, explanations, theorizing, and treatment suggestions of participants.
- In light of the lack of evidence for a causal relationship within the literature and the controversies that exist among researchers, we have tried not to pre-suppose a link between violence and eating disorders. Rather, our conclusions are based on the published literature and on data gathered from the service providers. We have placed the perspectives of the service providers at the centre of the paper in offering an explanatory framework for the links observed.
- We recognize that the diversity of opinion within the research is echoed among both the service providers and the advisory panel members who participated in this project. We have attempted to strike a balance in negotiating between those who cautioned us not to overstate the connection between violence and disordered eating and those who feared that we would understate the link. We have attempted to best represent the overall input of our sample.
- We have placed a high emphasis on the confidentiality of our participants, recognizing that many of the service providers consulted may know each other. Because this report is intended for distribution in a wide variety of contexts, we have not highlighted or analyzed responses by location, type of service provided, or other potentially identifying information. We acknowledge, however, that this would be another set of informative variables in understanding an abuse-eating disorder connection and in understanding the service provider relationship to that connection.
- Following the tradition of other published qualitative studies (cf. Websdale, 1995a; Websdale, 1995b; Websdale, 1998), the literature and the responses of the service providers have been integrated throughout the report rather than separated. This approach was selected in order to allow the reader a more direct comparison between the research and our “field data.” The literature was also used to provide context and to develop points raised by participants.

- This study was based on data collected from professionals and frontline workers in community, clinical, and private practice settings. We recognize that our participants use a wide scope of definitions and diagnostic criteria. We expect that this sampling represents the variety of perspectives and services being offered across Canada.
- One outcome of this review was a call by service providers for the integrated treatment of eating disorders and violence. This report therefore ends with a proposed integrated treatment model, emerging out of those factors identified as critical by Canadian service providers. Members of the advisory panel reviewed the model at the November 1999 teleconference.

THE NATURE OF VIOLENCE AGAINST WOMEN AND GIRLS AND OF EATING DISORDERS IN CANADA

PREVALENCE

Both survivors of family violence and those experiencing eating disorders in Canada are predominantly girls and women. Over 90% of individuals with anorexia and bulimia are female (Kuba & Hanchey, 1991; Prince, 1985; Siever, 1996). "Few disorders in general medicine or psychiatry are as skewed in gender distribution as eating disorders" (Andersen & Holman, 1997: 391). Canadian service providers working in the area of violence and/or eating disorders reported that the overwhelming majority of clients, callers, and workshop participants are girls and women. In fact, the proportion of female service users is generally over 90%. In addition, between 80 and 99% of those who experience *both* violence and eating disorders are female. Evidence suggests that the prevalence rates for eating disorders among girls and women (American Psychiatric Association, 1994; Hsu & Zimmer, 1988),² domestic violence against women (BC Institute Against Family Violence, 2001), and spousal killings of women (Duffy, 1998) are currently rising.

Violence Against Women and Girls

Approximately 80% of First Nations women have experienced physical, sexual, or emotional abuse (Duffy, 1998; Frank, 1992). Aboriginal women face 8 times the risk of battering as do non-Aboriginal women in Canada (Duffy, 1998). A third of First Nations women, on average, are abused by their partners (Frank, 1992), and between 75% to 90% of women living in some northern Aboriginal communities are battered (Dumont-Smith & Sioui Labelle, 1991). "Aboriginal people emphasize that family violence is not a tradition. Rather, family violence has become a problem following impacts of colonization" (Frank, 1992: 7). Another form of post-colonization violence experienced by First Nations families has been the application of child apprehension and adoption policies that have disproportionately placed children into non-First Nations families (Jiwani, 1998). Over 52% of all apprehended children in British Columbia are Aboriginal and 78% of Aboriginal children in permanent care in B.C. are placed in non-Aboriginal homes or facilities (Fournier & Crey, 1997). Through this process many children have experienced racism, dislocation, the erasure of cultural identities, and the severance of kinship, community, and ancestral ties (MacDonald, 1985). This is compounded by findings that 43% of Aboriginal and non-Aboriginal foster children are subjected to violence within the foster home setting (Kufeldt, Baker, Bennett & Tite, 1998). Within foster homes, 72% of girls report strong feelings of sadness and 18% of girls participate in deliberate self-injury (Kufeldt et al., 1998).

Over 40% of women with disabilities have been or are in abusive relationships (DAWN, 1989). Abuse experienced by women with disabilities is four times the national

² Note that Hof and Nicolson (1996) have argued that the apparent increase in eating disorders may both reflect changing norms of thinness and increased psychiatric labeling of women who defy feminine script.

average (Razack, 1994) and is mostly inflicted by family members or caretakers. Of women with disabilities surveyed by the DisAbled Women's Network, 37% reported abuse by parents and 17% reported abuse by spouses (DAWN, 1989). Two-thirds of able-bodied and disabled older adults who face abuse are women (The Senior Women Against Abuse Collective, 1989). Nearly 80% of this violence occurs at the hands of family members (Grandmaison, 1988).

Between 10% and 25% of non-Aboriginal, able-bodied Canadian women have experienced violence from their husbands or common-law spouses (Benson, 1995; Duffy, 1998; Statistics Canada, 1993). Reported rates are highest in B.C., where 1 in 3 women is assaulted by her spouse (BC Ministry of Health and Ministry Responsible for Seniors, 1995; Statistics Canada, 1993, 1995). About 8% of adult women in Canada have been sexually assaulted by past or current spouses (Duffy, 1998). Approximately 20% of lesbians in Toronto report being battered by a previous or current partner (Ristock, 1991). Children witness abuse between spouses or partners in 40% to 100% of violent relationships (Alliance of Five Research Centres on Violence, 1999; Statistics Canada, 1993).

Among non-Aboriginal, able-bodied girls, 12.5% (1 in 8) are seriously sexually abused and 20% (1 in 5) are physically abused (Duffy, 1998). In addition, a quarter of girls attending secondary school experience sexual or physical assault by a boyfriend (Duffy, 1998).

Eating Disorders

Overall prevalence of eating disorders

The Canadian Pediatric Society (1998) has reported that eating disorders are the third most common chronic illness among adolescent females. Over 65,000 young women in Québec between 14 and 25 years old are affected by eating disorders each year (Clinique St-Amour, 2001). Over 70,000 women meet clinical criteria in Ontario, a prevalence rate of 1 to 2% of the general population (Kraft, 1998). The American Psychiatric Association documents the prevalence rate of eating disorders as 1 to 4% (Joiner & Kashubeck, 1996). People(s) of colour(s) represent 1.8% to 5% of referrals for anorexia and bulimia in the U.S. (Davis & Yager, 1992).

Prevalence rates for bulimia

Epidemiological data from the Mental Health Supplement of the Ontario Health Survey suggest that the prevalence rate of bulimia for Ontarians over 15 years of age is 0.5% (Offord, Boyle, Campbell, Goering, Lin, Wong, & Racine, 1996). In a review by Woodside (1995), bulimia nervosa was estimated to affect 1 to 1.5% of women. This same clinician reported that binge eating in the absence of purging is much more common. In interview-based studies with rigorous diagnostic criteria, Garfinkel et al. (1995) found lifetime prevalence rates of 1.6% to 2.8% for Canadian women. The Anorexia and Bulimia Nervosa Foundation of Victoria (2000) estimates that the prevalence of bulimia in post-secondary students, however, may be as high as 1 in 6.

Epidemiological data from U.S. prevalence studies suggests that approximately 1.0% (1000 per 100,000) of girls demonstrate bulimia nervosa (Hoek, 1995). This estimate is consistent with findings of other researchers who report that 1 to 3% (leGrange, Telch, & Agras, 1997) and 1 to 2% (Pike & Walsh, 1996) of western women are affected.

Prevalence rates for anorexia

Garfinkel et al. (1995), of the Clarke Institute of Psychiatry in Toronto, observe that full-syndrome anorexia nervosa affects 0.56% of the population and that partial-syndrome anorexia nervosa is present in 1.4%. The Canadian Pediatric Society has documented an increase in eating disorders over the past 30 years and reports that they now affect up to 5% of adolescent women. Clinique St-Amour (2001) reports that about 1 to 2% of young adolescents will develop anorexia. Hoek (1995) has reviewed epidemiological data and reported average prevalence rates of anorexia nervosa of 280 per 100,000 (0.28%) among U.S. girls.

Breaking prevalence rates down

According to the Anorexia and Bulimia Nervosa Foundation of Victoria (2000), "recent findings indicate that both disorders affect all sections of the community and any type of family." Anorexia and bulimia impact women across race and culture (Bryant-Waugh & Lask, 1991; Daniels, 2001; Davis & Yager, 1992; DeAngelis, 1997; Field, Colditz & Peterson, 1997; Ford, 1992; Joiner & Kashubeck, 1996; leGrange, Telch & Tibbs, 1998; Pike & Walsh, 1996; Prince, 1985; Root, 1990), class (Gard & Freeman, 1996), and sexuality (Heffernan, 1996).

No significant differences have been found between women of colour and white women in reports of eating patterns or in the prevalence of eating disorder diagnoses (leGrange et al., 1997). The largest comprehensive study to date comparing black and white women, conducted by the National Institute of Health, has demonstrated roughly equivalent rates of bulimia and binge-eating (DeAngelis, 1997). In another study, African American women were more likely than white, Asian, and Latina women to have induced vomiting during the last month (Field, Colditz, & Peterson, 1997). High levels of disordered eating attitudes and behaviours have also been reported among adolescent Mexican-American women (Joiner & Kashubeck, 1996).

Within U.S. samples, 11% of Pueblo and Latina/o high school students (86% female) met *DSM-III* criteria for bulimia (Smith & Krejci, 1991). Higher rates of eating disorder symptoms have been found for Aboriginal girls and women (Crago, Shisslak & Estes, 1996; Rosen, Shafer, Drummer, Cross, Deuman & Maimberg, 1988; Smith & Krejci, 1991; Snow & Harris, 1989). In particular, Aboriginal adolescents have demonstrated higher rates of self-induced vomiting and binge eating (Smith & Krejci, 1991), depression after bingeing (Snow & Harris, 1989), and high levels of body dissatisfaction and fear of weight gain (Smith & Krejci, 1991; Snow & Harris, 1989).

Research in the U.S. has shown that 0.49% of lesbians currently meet diagnostic criteria for anorexia, and that 4.9% of lesbians have had anorexia in the past (Heffernan, 1996). This same research has demonstrated that 0.98% of U.S. lesbians meet diagnostic standards for bulimia nervosa.

Diagnostic criteria

A substantially larger proportion of women demonstrate bulimic symptomatology but do not meet criteria required for diagnosis of bulimia nervosa. Both service providers and members of the advisory panel cautioned that prevalence rates of anorexia and bulimia are unable to capture the broader continuum of “troubled eating” that girls and women experience. One advisor noted that diagnostic rates “certainly tend to ignore the concerns of individuals with food and weight issues that fall outside the very rigorous criteria for *DSM* classifications.” Another indicated that “there’s ample evidence to suggest that even amongst junior high and high school girls, many ... have sub-clinical eating disorders [and] may never reach the statistical threshold for falling into this group.” Estimates from the literature suggest that up to 19% of female students report bulimic symptoms in the absence of full clinical bulimia nervosa (Hoek, 1995; Zerbe, 1992). Approximately 5% of lesbians demonstrate binge eating disorder without meeting full diagnostic criteria for bulimia nervosa (Heffernan, 1996). One of our advisory committee members speculated:

... that you see a number of women who are victims of violence and they may have eating disorders, but they may not have any disorders that put them at medical risk. And if they don’t have eating disorders that compromise them medically, however you frame it, there just isn’t anything out there for them.

While the woman may be experiencing disordered eating, she may not be clinically diagnosable as having an eating disorder. Another reported:

As a front line worker, I work in an organization where I’m required to go by the *DSM-IV* criteria, and if my client hasn’t missed her periods for a sufficient amount of time, then supposedly she’s not anorexic which is kind of ridiculous with regard to providing service, given you want to do it in as timely a manner as possible.

Garfinkel et al. (1996b) have concluded that prevalence statistics based on diagnostic criteria are informative in a clinical setting but do not take into account the continuum of vulnerability to disordered eating. To work effectively with individuals experiencing high risk for disordered eating, all concerns regarding weight and shape issues and troubled relationships to food must be considered. Such difficulties may develop into more serious eating disorders or may still impact the psychological and physical functioning of the individual.

IMPACT

Both family violence and eating disorders are associated with serious physical and psychological health impacts. Many of these outcomes are common to both a history of abuse and to disordered eating. In fact, this similarity has been one factor that has led researchers to investigate the possibility of a connection. Anorexia, bulimia, and a history of abuse have all been linked to:

- feelings of shame and guilt, low self-esteem, a sense of inadequacy, and negative attitudes toward the body (Brown, 1997; DeGroot, Kennedy, Rodin & McVey, 1992; Herzog, Staley, Carmody, Robbins & van der Kolk, 1993; Schaaf & McCanne, 1994; Welch & Fairburn, 1996)
- problems with intimacy and trust (Brown, 1997; Herzog et al., 1993)
- negative feelings about sex (Brown, 1997)
- a sense of powerlessness in relationships (Schaaf & McCanne, 1994)
- a greater risk for alcoholism and substance abuse (Everill & Waller, 1995b; Schaaf & McCanne, 1994)
- depression and post-traumatic stress disorder (Everill & Waller, 1995b)
- self-harm through cutting, burning, scratching, and bruising (Brown, 1997; DeGroot et al., 1992; Everill & Waller, 1995b).

Within both eating disorders and family violence, the body is the physical site where issues of power and control are expressed. Through abuse, this power is stripped; through anorexia and bulimia, control is re-asserted. But both disordered eating and abuse are often shrouded in secrecy and stigma. As a result, many girls and women remain at risk of serious medical complications, injury, and psychological impact.

GIRLS AND WOMEN DYING

Between 1974 and 1993, approximately 75 women each year were killed by their spouses in Canada (Duffy, 1998). Women are 9 times more likely to be killed by spouses than by strangers (Duffy, 1998). Aboriginal women are 6 times more likely to be killed by a spouse than are non-Aboriginal women (Duffy, 1998). About 43% of all wife assaults result in medical attention and a third of women living in violent relationships fear for their lives due to the severity of the violence (Statistics Canada, 1993). In fact, fear of injury or death is as high for women who are raped by their husbands and their dates as it is for survivors of stranger rapes (Koss, 1993). In addition to the threat posed during abuse, women who have endured childhood sexual abuse, adult sexual assault, or battering are more likely to attempt suicide (DeGroot et al., 1992; Schaaf & McCanne, 1994; Yoder, 1999). Suicide is an even greater concern for high-risk groups, such as Aboriginal and/or lesbian, bisexual, or Two-Spirited³ youth and women.

³ "The term 'Two-Spirited' originates from the First Nations recognition of the traditions and sacredness of people who maintain a balance by housing both the male and female spirit" (Deschamps and Wahsquonaikzhik, 1998: 10).

Approximately 90% of individuals who die from anorexia and bulimia are girls and women (Gagnon, 1996). Eating disorders are the most life-threatening of all psychiatric conditions (Zerbe, 1992). About 5% of girls and women with anorexia die from complications or starvation within the first 5 to 8 years (Woodside, 1995). Risk of mortality increases each year. Over a 20-year period, 13% to 20% of women with anorexia will have died (Woodside, 1995). Over a 3 to 5 year period, there is a 5% fatality rate for bulimia (Woodside, 1995). Adolescent women comprise 20% of the deaths from anorexia (Gagnon, 1996). Severe medical illnesses accompany chronic anorexia in 56% of cases (Herzog, Deter, Fiehn & Petzold, 1997). Many health effects can be irreversible, even after recovery (Canadian Pediatric Society, 1998). Health outcomes for older women with eating disorders are particularly poor (Cosford & Arnold, 1992), and overwhelmingly lead to death in women who are 70 years of age or older (Gagnon, 1996).

While many outcomes associated with family violence are serious, anorexia and bulimia are health and life threatening. Both abuse and disordered eating can be debilitating to the woman and potentially result in her death. Although the shared serious nature of family violence and eating disorders does not indicate their correlation, it does suggest that their co-occurrence may have critical consequences. Moreover, the impact of eating disorders and of violence extends beyond the individual woman and girl within whom they have been embodied. These issues also impact on the health and the social and economic fabric of the broader community in which they are embedded.

LINKS BETWEEN VIOLENCE AGAINST WOMEN AND GIRLS, ANOREXIA, AND BULIMIA

Nearly two-thirds (63%) of the 123 service providers who responded reported seeing a link among their clients between family violence and disordered eating. Only 6.5% said that there is no link. Estimates from Canadian service providers regarding the number of women and girls with anorexia and bulimia who have experienced family violence varied widely. The average estimate across service providers was that 50% of the women they treat for disordered eating have a history of abuse of some form – sexual, physical, emotional, or psychological.⁴ The average estimates were lower (30%) for Québec and higher (60%) for Saskatchewan and Nunavut/NWT.

About 9% of those who identified a link indicated that it was a moderate or weak one. Approximately 8% of informants emphasized that violence was a contributing factor but not the only cause. Nearly half (46%) of service providers contacted, however, characterized the connection between violence and eating disorders as “definite,” “strong,” “absolute,” “always,” “usually,” or “almost always.”




Figure 1: Responses by Canadian Service Providers to a Possible Link between Family Violence and Eating Disorders

⁴ The average estimation figure of 50% represents the mean, median, and the mode of the responses provided to us by the 123 service providers across Canada. Standard deviation of the responses was 25.14%.

As with the published literature, there was a high variability among responses from service providers, reflecting differing definitions of abuse, diagnostic criteria for eating disorders, the type of population served, and the type of treatment provided. The subsequent sections of this report explore further our respondents' experiences with varying forms of abuse, different age groups, observations about mediating factors between abuse and disordered eating, explanations for their experiences, and approaches to treatment.

ABUSE OF CHILDREN AND THE DEVELOPMENT OF EATING DISORDERS

Childhood Sexual Abuse

Sexual abuse has been a topic of research and interest since the mid 1980s (Thompson, 1992). With this growing interest has come an exploration of childhood sexual abuse as a possible contributing factor to anorexia and bulimia. Although estimates range widely, published research indicates that, overall, about 30% of women with eating disorders have been sexually abused (Connors & Morse, 1992). Data collected on 25,838 British Columbian adolescents found that 28% of young women reporting problem weight control practices also reported sexual abuse (McCreary Centre Society, 1998). Nearly a quarter (23%) of Canadian service providers that we contacted reported a clear link between sexual abuse during childhood and the development of anorexia or bulimia.

The connection between sexual abuse and eating disorders may be influenced by a number of factors. Women are at greater risk for eating disorders if they have had higher numbers of unwanted sexual experiences (Mullen et al., 1993), higher numbers of abusers (Davenport et al., 1994), or if physical abuse has accompanied sexual abuse (Miller, 1996; Wonderlich, Brewerton, Jolic, Dansky & Abbott, 1997).

Women who have experienced past extrafamilial child sexual abuse (i.e., abuse by someone outside the family) have reported bingeing, vomiting, fasting, and using diet pills more frequently than those who have not (Hernandez, 1995). Intrafamilial (abuse by a family member) sexual abuse (Baldo, Bostwick, Wallace & O'Halloran, 1996; Waller et al., 1993), particularly by a close male relative (Mullen et al., 1993), has been linked to greater likelihood and more serious manifestations of eating disorders. Canadian researchers have also observed a difference in the type of eating disorder that develops. For example, women with bulimia who purge may be more likely to have experienced sexual abuse than

women with bulimia who do not purge (Garfinkel et al., 1996a). High rates of sexual abuse have been found in Canadian women for both types of anorexia (Garfinkel et al., 1996b).

"It is a chilling reality for a staggering number of girls that sexual activity is equated with violence and violation long before they are introduced to life-affirming sexual activity based on mutual consent."

- Thompson (1994)

Childhood Physical Abuse

The role of childhood physical abuse in the development of disordered eating has received much less attention than has childhood sexual abuse (Reto et al., 1996; Rorty & Yager, 1996). There is, however, some evidence to support a connection. Over a third of British Columbian adolescent women surveyed who reported frequent binge eating or vomiting, frequent use of diet pills, or frequent weight loss efforts reported physical abuse histories (McCreary Centre Society, 1998). Aboriginal, African-American, Latina, and white U.S. children, adolescents, and women who have been physically abused have reported higher incidences of eating disorders and more severe cases of bulimia (Hernandez, 1995; Reto et al., 1996). Women with bulimia are more likely to have experienced repeated and severe physical abuse and to have been abused within the year preceding the onset of symptoms (Welch, Doll & Fairburn, 1997; Welch & Fairburn, 1996). Moreover, physically abused women may have more difficulty identifying hunger and fullness than sexually abused women (Schaaf & McCanne, 1994). Bodily shame (Andrews, 1997) and an emotionally disturbed family environment (Schmidt, Humfress, & Treasure, 1997) have been found to further strengthen links between physical abuse and bulimia. In particular, research has suggested that bulimia may offer a dissociative response to physical abuse and the feelings of shame that derive from it (Nagata, Kiriike, Kawarada, & Tanaka, 1999; Reto, 1998).

Of the Canadian service providers we contacted, 1 in 7 (14%) said that there was a link between physical abuse experienced during childhood and the development of eating disorders. One B.C. informant estimated that about 60% of eating disorders are directly tied to physical abuse. Taken together, the published research and the experience of service providers indicate that a history of physical abuse may play a contributory role in the development of eating disorders, particularly bulimia.

Family Interactions during Childhood

Seven of the service providers (5.7%) emphasized family dysfunction and poor family relations as factors in the development of anorexia and bulimia. In particular, they identified poor communication, lack of conflict resolution, higher levels of conflict, perfectionist expectations, and witnessing of woman or child abuse. One service provider defined what she termed “intellectual abuse” as a form of verbal abuse by middle-class and affluent parents in which sarcasm, perfectionism, rigid control, and high expectations undermine the child’s self-esteem. Factors that have been highlighted in the published literature include parental indifference, excessive parental control, family discord, and parental disapproval (Schmidt et al., 1993). Leighton (cited in BWSS, 1997) also notes that the witnessing of woman abuse by children is linked to the development of eating disorders.

An additional 14% of the service providers said that *emotional or verbal abuse* during childhood was connected to the emergence of anorexia and bulimia. One informant observed that a background of verbal abuse was very common, and another suggested that

most clients experienced emotional rather than physical abuse. Thompson (1994) has observed that many eating disorders begin as a “search for refuge” from physical abuse or such forms of emotional abuse as verbal insults, accusations, refusals to give reassurance of love, neglect of basic needs, and lack of physical touch. Rorty and Yager (1996) have pointed out that a continuum of maltreatment must be considered that includes parental/caretaker intrusiveness (e.g., opening mail, eavesdropping, reading diaries), and sexualized relationships (e.g., genital exposure, sharing pornography, etc.).

Service providers also pointed to damaging messages about food that children receive within the family, including verbal abuse about size and fatness, withholding of food, forcing children to eat food, and social stigma attached to types and quantities of food. Mealtimes may be stressful and may become an area in which parental authority, control, and abuse are exerted (Miller et al., 1993a). Women with bulimia have commented that “‘it was a relief when my father wasn’t at dinner’; ‘my father commented about my mother’s weight when I was young’; ‘when I was hurt or upset my mother would offer me food as a special treat’; ‘both my father and mother made me eat food I didn’t like’” (Miller et al., 1993a: 311). If mealtime control is combined with patterns of sexual and physical abuse, the food itself may become a symbol of conflict, intrusion, and pain. Women with bulimia have reported more negative mealtime and food-related experiences, high levels of stress and conflict during meals, parental use of food as a tool for punishment or manipulation, and a family emphasis on dieting and weight (Miller et al., 1993b). Family interactions during meals and around food can be especially pertinent to the maturing girl if access to food is gendered. In some homes, boys may be given more food and encouraged to eat while girls entering adolescence are increasingly pressured to reduce food consumption in order to attract a male partner (Thompson, 1992).

YOUTH AND ADOLESCENT EXPERIENCES OF VIOLENCE, ANOREXIA, AND BULIMIA

The connection between family sexual, physical, and emotional violence and the development of disordered eating in adolescents is often included in research addressing childhood experiences. But there are also types of abuse that are uniquely experienced by adolescents.

In an epidemiological study examining adult abuse and *DSM-III-R* disorders, Danielson et al. (1998) reported that between 35 to 50% of young adults were involved in some level of physical partner abuse. Mercer (1987)⁵ found that among the Toronto secondary school women he surveyed, 20% reported experiencing abuse in their romantic relationships. The Nova Scotia Advisory Council on the Status of Women found that among young women in dating relationships, 11% experienced sexual abuse, 32% reported emotional abuse, and 18% faced physical abuse (Day, 1990). Moreover, adolescent wives (ages 15 to 19) are murdered three times more often than adult wives (Canadian Centre for

⁵ Research by Mercer (1987), Canadian Centre for Justice Statistics (1994), and Leighton was integrated and reported by Battered Women’s Support Services (1997).

Justice Statistics, 1994). Unfortunately, there has been virtually no research to examine the role of battering in adolescent girls' romantic relationships and their pre-existing manifestation or subsequent development of eating problems. One of the service providers, however, informed us that girls with bulimia and anorexia talk a lot during group sessions about violence in schools, with friends, or in dating relationships. In addition, whereas boys under the age of 13 are more likely to be physically abused in the parental home, girls are more likely to be physically abused when older (Gadd, 1997).

Lesbian and gay youth are confronted with homophobia both within the family and from peers. Adolescence is a time when they face pressures to heterosexually date and to engage in heterosexual discourses, as well as to make decisions about "coming out." Between 20 to 50% of all street youth in Canada are lesbian or gay (Deschamps, 1998), many having left to escape homophobia within the family and within peer environments. In his thesis research, Samis (1995) found that 5% of lesbians and 11% of gay men in the Vancouver region reported having been gay-bashed by family members. Thompson (1994) has highlighted the role of homophobia as a form of "psychic violence" in the development of eating disorders. Lesbians she interviewed reported bingeing in response to the isolation of coming out (Thompson, 1992). Research in British Columbia has found that 15% of non-heterosexual youth report purging at least once in a while compared to 6% of heterosexual youth (McCreary Centre Society, 1999). One service provider noted "many instances" of eating disorders where the women consulting her "have questioned their sexuality." The connection between homophobia and eating disorders is further supported by the greater prevalence of anorexia among young gay men than among young heterosexual men (French, Story, Remafedi, Resnick & Blum, 1996; this point was also raised by one of the service providers we consulted).

Adolescence is accompanied by a number of lifestyle changes: assertion of independence, part-time jobs, spending more time away from home, driving, increased body fat, occupation options, increased importance of peers, and social pressure regarding appearance (Child and Family Canada, 1994). The adolescent woman may encounter prohibitions on freedom and exertions of parental control as she seeks her adulthood (cf., Hill & Holmbeck, 1987). She receives messages that her sexuality must be protected, and experiences the gender-specific nature of this standard if her brothers have later curfews, less surveillance, and fewer restrictions (Atwood, 2001; Peters, 1994; Zani, 1991). In this way, she increasingly becomes the body that cannot be trusted (cf., Gremillion, 1992). This situation can be compounded when sexual remarks and jokes are made about her body by family members or peers, when her developing secondary sexual characteristics become the object of harassment, or if she has or is experiencing sexual abuse (cf., LaBarbera, 1984; Larkin & Popaleni, 1994; Smith, 1997). Young women may be encouraged by family to develop goals and body images that are conducive to upward class mobility. In fact, some researchers have hypothesized that upward mobility is a contributing factor to eating disorders. Weight loss has become a symbol of interpersonal and financial success (Striegel-Moore, 1995). Thinness in North America represents competence, intelligence, assertiveness, and self-control (Smith, Waldorf & Trembath, 1990). Women are often required to modify their bodies for obtaining access to employment, education, and

economic resources (Littlewood, 1995). Young women often face pressures from parents toward thinness, dieting, and lighter or restricted eating (Tsiantis & King, 2001; Vincent & McCabe, 2000). The degree of family pressure to control weight has been found to moderate the relationships among thinness norms, body dissatisfaction, and disordered eating (Twamley & Davis, 1999). More serious manifestations of eating disorders have been linked to more critical maternal attitudes toward weight during adolescence (Ritter, 1998). Strong family relationships, in contrast, have been found to decrease the risk for eating disorders among abused youth (Neumark-Sztainer, Story, Hannan, Beuhring, & Resnick, 2000). Consistent with this finding, research suggests that youth who have spent more time in foster or group homes (Garfinkel et al., 1995), and homeless youth (Freeman & Gard, 1994) are at greater risk for bulimia.

Both race and class assimilations are tied to thinness and dieting (Root, 1990). In interviews conducted by Thompson (1992), African-American and Puerto-Rican women both highlighted their experiences of family pressure to emulate white, middle-class standards of thinness. For example, one woman explained that as her family class shifted from a working class to an executive, upper-class level, her family moved from big meals, chubby children and keeping plenty of food in the house to an insistence on elegance, thinness and putting her on diet pills. Root (1990) further discusses how the devaluation of non-European cultures causes adolescent women of colour to move toward European values of beauty and reject their cultures of origin. She warns that this process converges with the developmental point at which women are most vulnerable to eating disorders. One African-American woman, for example, related her eating disorder to statements by her white grandmother that she would never be as pretty as her cousins because of their lighter skin (Thompson, 1992). Racialized identity is further impacted by the presence of abuse. "Sexual violence shakes up what 'home' means. Since racial and cultural identity is primarily taught in the home, this socialization process is inevitably disrupted when the home is no longer a refuge, but rather a place of stress and fear" (Thompson, 1994: 68). One service provider added that the negotiation for young first generation Canadian women between the dominating culture and the struggle of their parents to preserve cultural identity is one precipitating factor for eating disorders.

Family dynamics may have specific impact for adolescents with chronic illness and disability. Youth with chronic illness have been found to be at greater risk of disordered eating than those without (Neumark-Sztainer, Story, Falkner, Beuhring & Resnick, 1998). This same sample of youth also experienced lower levels of family communication, parental caring, and parental expectations. In addition, adolescents with chronic illnesses have reported more sexual and physical abuse (Neumark-Sztainer et al., 1998). Such research suggests that an interaction of non-supportive family dynamics, family violence, and the presence of chronic illness may put youth at greater risk of developing anorexia or bulimia. Women with disabilities indicate confronting similar pressures as able-bodied women to conform to standards of body shape, size, and weight (McCarthy, 1998; Watson, 1999). But, research on samples of women with arthritis, visible blood vessel conditions, skin conditions, diabetes (Ben-Tovim & Walker, 1995), spinal cord injury (Beatus, 1997), learning disabilities (McCarthy, 1998), and other visible physical disabilities (Watson,

1999) has generally reported that women with disabilities do not necessarily disparage their bodies more or demonstrate a more negative body image than able-bodied women. Rather, disabled women have indicated that, as a result of discrimination, they experience a stronger pressure to “dress for success” (Watson, 1999).

In addition to the above, informants who were consulted for this study noted the impact of harassment as experienced by young women in high school. Violence within the school environment was also emphasized. One service provider observed that 80% of girls in her practice with eating disorders experience some form of violence at school, and that experiences of harassment are more prevalent than are verbal, sexual, and physical abuse.⁶ Another informant mentioned an “increasing amount of emotional violence exchanged between the girls, which takes the form of verbal ‘put downs’ or undermining with malicious intent.” Teasing by peers has been found to predict both body esteem and eating behaviour among adolescent girls (Lieberman, Gauvin, Bukowski, & White, 2001). Indeed, recent research into the role of peers in disordered eating has suggested that members of the same friendship cliques tend to share similar body image concerns and dieting behaviours (Paxton, Schutz, Wertheim, & Muir, 1999; Stump, 1995). Those girls whose friends participate in extreme weight-loss behaviours are more likely to engage in these same behaviours (Paxton et al., 1999). Even in third and fifth grades, children’s eating and body concerns have correlated to being liked by peers (Oliver & Thelen, 1996).

VIOLENCE AND EATING DISORDERS IN ADULTHOOD AND AGING

Battering

There have been few studies examining the relationship between eating disorders and physical abuse experienced during adulthood (Kaner, Bulik & Sullivan, 1993). Some information has come from research done in the U.S. In a study conducted about a decade ago, 45% of women with bulimia reported being physically victimized at least once during an adult relationship (Root & Fallon, 1988). Of these, 23% had been raped, 23% had been battered, and 6% had been both raped and battered. In a more recent U.S. study, almost two-thirds of women interviewed who experienced severe spousal abuse met the criteria for one or more psychiatric diagnostic category and had elevated rates of mood, anorexic/bulimic, and substance use problems (Danielson et al., 1998). Perhaps the most informative study is that by Kaner et al. (1993). These researchers found that 40% of women with bulimia compared to 5.9% of women without bulimia had been battered 1 to 3 times per week during an adult relationship. The risk of battering was 6.8 times higher for women with bulimia. Moreover, women with bulimia were more likely to blame themselves for the abuse and to feel that it was deserved than were those women without bulimia. An additional factor occurs if the woman leaves her batterer and is confronted with initial or greater poverty, in which food becomes one of the least expensive substances available as a method for coping (Thompson, 1992). The findings of these studies suggest that battering and other forms of abuse in women’s heterosexual and lesbian romantic relationships may be factors in the development and manifestation of eating disorders.

⁶ This observation is supported by the statistical profile of youth violence which demonstrates low levels of physical violence between girls or where girls are the perpetrators (Reitsma-Street, 1999).

In this study, four service providers linked anorexia and bulimia with the experience of physical assault. One informant reported that “the most common scenario is the married bulimic woman with a history of domestic violence,” noting, in addition, that children who come from violent homes are also at risk. A second service provider similarly observed that eating disorders tended to be manifest among women in their 30s and 40s who have a history of child abuse and more recent experiences of domestic violence. For facilities and practitioners who received referrals, some noted that many of these came from women’s shelters. Others reported that they sometimes found it necessary to refer women with eating disorders to women’s shelters because of the immediacy of the experience of abuse or because they lack knowledge in the area of abuse. Our advisory committee noted that people tend to:

... talk a lot about integrating treatment in terms of when women come to seek treatment for their eating disorder. What about when women come to seek support, emotional support or advocacy for the fact that they’re in an abusive relationship and eating disorders come up in the course of the counselling session or in the course of our work with her? Our focus seems to be the other way: when women go and get treatment for their eating disorder and then how do we integrate issues of violence into that?

Sexual Trauma in Adulthood

Evidence has been found by numerous researchers for an association between past sexual trauma and bulimia (Baldo, Bostwick & Baldo, 1996; Connors & Morse, 1993; Dansky et al., 1997; Everill & Waller, 1995a, 1995b; Lanzi, Balottin, Brisone, Citterio et al., 1997). In her study of African-American, Latina, Jewish, and white North American women, Thompson (1992) found that sexual abuse history was the trauma that participants most frequently related to the development of their eating problems. In a national U.S.-based study of 3006 women, participants with bulimia had experienced higher rates of rape, sexual molestation, aggravated assault, direct victimization, and current and lifetime post-traumatic stress syndrome (Dansky et al., 1997). In a college sample of women, just over twice as many women with bulimia (56.8%) as women without bulimia (26%) reported having narrowly missed being sexually assaulted (Beckman & Burns, 1990). Both rape and oral sexual abuse have been found to be related to bulimia (Welch & Fairburn, 1996). Canadian research has also shown that 82% of federally incarcerated women and 72% of provincially incarcerated women have experienced either sexual or physical abuse or both. Within this same population, 59% disclosed that self-injurious behaviour and disordering eating were common (National Crime Prevention Council of Canada, 1995).

Six (5%) service providers we consulted reported that sexual assault and rape were linked to disordered eating in Canadian women. One informant further emphasized the role of sexual harassment.

Aging

Between 1986 and 1988, elder abuse in Canada increased by almost 20% (National Aging Resource Centre on Elder Abuse, 1990). Overall, elder abuse has been estimated to affect 1% to 10% of the senior population, with incidences as high as 36% for physical abuse and 81% for psychological abuse within some institutions (Patterson, 1993). Approximately 4% of Canadian elders living in private dwellings have reported being abused and a fifth of these have reported more than one type of abuse (National Clearinghouse on Family Violence, 1993). Older women are roughly one and a half times as likely to be abused as older men (5:3) (National Clearinghouse on Family Violence, 1993). In 1992, approximately 1.4 million women between 45 and 64 years of age were physically abused by spouses (Mother's Report Call to Action Violence against Midlife and Older Women, May 1994). In an Ontario survey, 20% of nurses and nursing assistants indicated that they had witnessed abuse of patients within nursing homes; 10% reported that other staff had hit or shoved patients; and 28% observed yelling, swearing, and embarrassing comments (National Clearinghouse on Family Violence, 2000a). Elder abuse may include beatings, sexual assault, physical restraint, failure to provide care or necessities, humiliation, isolation, intimidation, abandonment, and financial control. Moreover, seniors face increasing isolation through the death and loss of friends, siblings, and spouses or lovers. They may not have a network to seek help or shelter from abusive environments. Aging women are at particular risk of food deprivation and control if they are dependent on others to obtain, prepare, plan, and/or feed them meals. They may also face infantilization in their changing relationships with adults younger than them, which is often highlighted in interactions around feeding. For the older woman who is experiencing physical or sexual abuse, restraint, condescension, or neglect, it is the aging body that is the site of trauma. She may decide that it is simply "her time" and engage in reduced food consumption. She may alternatively seek control of her body again through food refusal or binge-purge cycles. Eating may be one of the few avenues through which the aging woman feels she may exert control. Wiederman (1996), for example, discusses an 86-year-old woman who, upon institutionalization, developed an eating disorder. Her own description of her environment highlighted noise and intrusion, a lack of control over food selection and preparation, frustration over a loss of independence, and poor quality food. For another 80-year-old white widowed woman, dieting and purging represented weight loss and relief that had been denied to her by her husband when he forbade her to seek a breast reduction (Beck, Casper & Andersen, 1996). She had continued to live with social and physical discomfort until the age of 73, when her husband died.

Eating disorders among post-menopausal women are under-diagnosed, not recognized as legitimate diagnoses, or met with skepticism (Cosford & Arnold, 1992). In one study of 5 patients over 55 years with eating disorders, it was found that all had been under the care of a physician, 2 were seeing a psychiatrist, and none had been diagnosed (Hsu & Zimmer, 1988). Yet, eating disorders are overwhelmingly fatal for seniors. Just over 78% of deaths from anorexia occur in individuals over 45 years of age (Gagnon, 1996). When older adults depend on family or institutions for finances, transportation, housing, or the provision of basic necessities, then they are at risk of abuse. If food is used

as a weapon in abuse or is provided in an atmosphere of dependence by the abusing individual, then eating may become an aversive experience that the older adult resists.

THE ROLE OF DISCLOSURE

Eighteen (15%) of the service providers consulted raised the issue of disclosure in assessing the link between violence against women or girls and eating disorders. Girls or women who are hospitalized or who seek treatment for anorexia or bulimia may be reluctant to disclose current or past abuse. Violence may be difficult to address due to the secrecy that surrounds abuse, the ongoing threat of the situation, and denial by girls, siblings, or parents that violence exists in their families. Secrecy may be exacerbated in rural or small town settings where lack of anonymity and stigmatization are concerns (Jiwani with Moore & Kachuk, 1998). One service provider also noted that "families with secrets like a history of violence are not likely to seek out this programme." Women and girls living with violence may be less apt or less able to pursue treatment for anorexia and bulimia. Moreover, because both the history of abuse or sexual trauma and having an eating disorder are stigmatized, the girl or woman who experiences both may be less likely to seek help or, alternatively, may seek help for one but be hesitant to disclose the other. As one advisory committee member commented, "This whole issue of accessibility and the fact that whether it's a history of violence, abuse, or an eating disorder, people are often very reluctant to disclose that. And the younger they are, the more reluctant they tend to be sometimes." It is important to note, however, that women and girls are more likely to reveal abuse if they are directly asked (National Clearinghouse on Family Violence, 2000b). Both the American College of Obstetricians and Gynecologists and the American Medical Association have recommended routine screening of women and girls for abuse. Nonetheless, across Canadian and U.S. studies, health care practitioners indicate that they do not routinely or directly ask about abuse (National Clearinghouse on Family Violence, 2000b).

One service provider suggested that the setting may be important to whether the woman or girl discloses abuse as a factor in her eating disorder. Several variables related to setting were discussed by informants. Some dietitians and nutritionists indicated that women were more likely to disclose to other types of mental health workers. In particular, it was suggested by one therapist that an examination of family systems and structures tended to encourage greater disclosures of violence. Three service providers indicated that group settings often led to higher disclosures and more discussions of abuse as a factor in eating disorders than did individual sessions. Abuse is then subsequently addressed in individual therapy, where at least one informant felt it was most appropriately addressed. On the other hand, one service provider emphasized that family violence was not usually raised in group sessions. It was also noted that when the connection between violence and disordered eating was directly addressed during the initial assessment, the frequency of disclosure was higher.

Higher levels of disordered eating have been found in women who have previously disclosed their abuse and received an adverse reaction (Everill & Waller, 1995a). Responses such as disbelief, blaming the girl or woman, ignoring, and punishment can further increase the woman's feelings of general worthlessness, inferiority, and stigmatization (Brown, 1997; Everill & Waller, 1995a). The woman may feel betrayed, distrust the experiences of her own body, or distrust her ability to communicate her experiences (Everill & Waller, 1995b). For the girl, disclosure may be followed by the removal of her father from the home and subsequent loss of income or scapegoating of her for "breaking up the family" (Friedrich, Urquiza & Beilke, 1986). In the context of ongoing discrimination in child apprehension, the Aboriginal woman may fear losing her children if she discloses eating disorders, alcoholism, or battering.

Like the service providers we interviewed, some researchers have suggested that the mixed research findings about the links between violence and eating disorders may be, in part, attributable to under-disclosure of abusive experiences due to denial, repression, and dissociation (Connors & Morse, 1993; Miller, 1996). This may particularly affect girls and women who have experienced more severe forms of abuse (Miller, 1996). Moreover, girls and women may not always label their experiences as abuse even when asked directly. For example, in one study, 42% of children who described multiple experiences of burning, biting, kicking, and beating with an object by parents did not self-label it as abuse (Reto et al., 1996). Miller (1996) has highlighted some strategies for assessment which tend to elicit greater disclosure. These include more inclusive definitions of abuse, the use of "inverted funnel questioning" (i.e., specific and overlapping questions about the experience), and interviewing. Detailed questionnaires combined with interview formats generate the most information (Connors & Morse, 1993). The use of interviewing is particularly important in light of one study in which a third (33%) of women who did not disclose child sexual abuse on an intake questionnaire later disclosed in a research interview (Miller, 1996). The validity of such interviewing is supported as reports of childhood abuse by women generally demonstrate high rates of corroboration, exceeding 75% (Weiner & Stephens, 1996). The National Clearinghouse on Family Violence (2000b) has recommended that women be asked about specific acts (e.g., slapping) rather than global problems (e.g., domestic violence), using either screening instruments (e.g., checklists) or private patient interviews.

AGE

Service providers presented a relatively consistent picture regarding ages at which women experience abuse, disclose abuse, and seek or require help. Overall, they indicated that the connection between violence and eating disorders affects women across the lifespan. While many eating disordered women have experienced abuse at an early age (for example, 9 to 10 years old), most of these women disclose the abuse and develop their eating problems when they are in late adolescence or early adulthood. Fewer women disclose current experiences of violence than discuss a history of past violence. One service provider observed that incest and sexual abuse are not always revealed by young women or

adolescent girls, but are discussed by older women who are more ready and who are no longer dependent on their families.

Women in their 60s and 70s were identified by service providers as being in an age group also suffering from eating disorders that are linked to abuse or violence. Research has suggested that childhood sexual abuse remains an important issue for later onset cases of eating disorders (Beck et al., 1996). Memories and associated emotions still persist as women age. Attaining closure on past experiences of abuse has been identified as an important part of recovery for older women with anorexia and bulimia (Beck et al., 1996).

Research studies examining violence against girls or women, anorexia, and bulimia are almost exclusively retrospective. Both adult women and adolescent girls are asked to recall abuse history and childhood environments (e.g., DeGroot et al., 1992; Miller, 1996; Moyer et al., 1997; Schmidt et al., 1993). Unfortunately, this approach tends to focus on adult samples remembering childhood through an adult perspective, rather than on adolescent samples at a time when many eating disorders develop (Herzog et al., 1991). The opportunity for youth to disclose both eating disorders and ongoing family or other forms of violence is critical to an integrated therapeutic approach, as well as to providing a better understanding of abuse as a contributing factor in the development of an eating disorder.

UNDERSTANDING THE LINKS

The explanations that service providers offered for a connection between abuse and disordered eating demonstrated five themes: causality; coping; control and power; self-esteem; and sociocultural context.

CAUSALITY: DO EXPERIENCES OF VIOLENCE LEAD TO EATING DISORDERS?

Many researchers and clinicians have observed high abuse rates among girls and women with anorexia and bulimia. However, some have questioned whether this connection simply reflects that both abuse and eating disorders occur predominantly among women and girls (Connors & Morse, 1993). If sexual abuse is a contributing factor to the development of eating disorders, then child sexual abuse must precede the onset of symptomatology. Some evidence exists that, at least for some women, sexual abuse does precede and contribute to their anorexia or bulimia. For over 90% of girls who have survived sexual abuse, the age of onset for eating disorders is *after* their first experiences of abuse (Herzog et al., 1993; Miller, 1996). Both research and one of the service providers have estimated the average age of first abuse for women with eating disorders to be about 9 to 10 years old (Miller, 1996). Approximately 10% of women who develop bulimia are known to have been sexually abused within a year preceding the onset of their symptoms (Welch et al., 1997).

Both researchers (Welch & Fairburn, 1996), and some of the service providers have cautioned against “oversimplifying” the relationship between family violence and eating disorders. “An approach insisting on a one-to-one or straightforward causal relation is also likely to miss many of the more individual and experiential aspects trauma plays for this particular population” (Brown, 1997: 195). Overall, both reviewers and many of the service providers have concluded that although child sexual abuse is neither necessary nor sufficient for the development of eating problems, it does act as a contributory factor (Connors & Morse, 1993; Dansky et al., 1997; Reto et al., 1996; Rorty & Yager, 1996). As one of our advisory committee members noted, “it’s not necessary or sufficient that people have violence in their background to develop anorexia or bulimia but there is a considerable link in considerable numbers of the cases.” A history of abuse was characterized as one contributor in a complex multideterminant model that includes both risk factors and resiliency factors.

COPING WITH VIOLENCE

Nine service providers told us that eating disorders are one type of coping mechanism for women and girls who have experienced violence. In particular, food is used to deal with stress and with more serious underlying family problems. Anorexia and bulimia may emerge as ways of coping with the pain or trauma from living with ongoing or past abuse.

Bingeing serves different, and potentially multiple, purposes for different individuals. For some women, eating is used as sedation, to alleviate anxiety, and to combat loneliness (Thompson, 1992). For adolescent women surviving sexual abuse, food may be the most accessible and socially acceptable drug available to them (Thompson, 1992). Dissociation may provide a connection between a history of sexual or physical abuse and subsequent eating disorders. Dissociation allows escape from trauma when physical escape is not possible (Everill & Waller, 1995a). In particular, bingeing can serve to block emotions, self-degradation, continued memories of the abuse, fear, self-blame, and anger from awareness (Everill & Waller, 1995b; Miller et al., 1993a; Wonderlich et al., 1997). Binge eating draws the girl's or woman's attention to the immediate present, thereby blocking depression, anxiety, and thoughts (Wonderlich et al., 1997). In addition, it induces a physiological alteration that is similar to the lasting chemical changes in the body that ensue following trauma (Wonderlich et al., 1997). In this way, it "anesthetizes" the woman, allowing her to numb feelings and attain relief (Brown, 1997; Thompson, 1992). Psychologically "leaving the body" during bingeing parallels "leaving the body" during sexual and physical abuse (Thompson, 1992). Moreover, the act of eating large amounts of food can encourage sleep and provide another way to block pain (Thompson, 1992). As the woman or girl blocks her feelings, memories, and experiences through bingeing, she may follow this with attempts to "cleanse" herself of violation through purging (Brown, 1997; Waller, Ruddock & Cureton, 1995). Thompson (1994) has described bulimia as a mechanism for "throwing up" abuse.

Childhood abuse can include starvation, force-feeding, forced ingestion, emotional abuse around eating and weight, forced eating of vomitus or spoiled foods, physical abuse during meals, use of food as rewards or bribes following abuse, and use of food in sexual acts (Brown, 1997). Within Canadian residential schools, food was often used as a weapon of abuse. Aboriginal children endured starvation (as punishment and as a cost-saving measure), food unfit for human consumption, the forced ingestion of vomit, inadequate nutrition for growth and subsistence, forced feeding, bread and water as punishment, and severe beatings for trying to acquire extra food or water (Chrisjohn & Young with Maraun, 1995). These food deprivations and punishments occurred in conjunction with extensive physical and sexual abuse. Research has suggested that young men are more likely to express pain resulting from abuse outward and young women are more likely to direct pain inward into self-injury and eating disorders (National Clearinghouse on Family Violence, 1997a). Moreover, when inflicted as abuse, food itself may become unappetizing, an aversion, or a provision over which the abused girl or woman is not accustomed to having control or access. Certain foods may trigger memories of ejaculation through their appearance or consistency, leading to their avoidance or purging (Brown, 1997). For girls and women who have histories of family or institutional violence, we are reminded that "their eating strategies began as logical solutions to problems rather than problems themselves as they tried to cope with a variety of traumas" (Thompson, 1992: 558).

CONTROL AND POWER

Increasingly, there is a shifting focus in the theoretical model used to understand disordered eating from a model in which a desire for thinness is regarded as the central factor in disordered eating to one that emphasizes control and powerlessness (Thompson, 1992, 1994). This shift has provided more clarity in understanding the connection between violence against girls and women, anorexia, and bulimia.

Work by Sandy Friedman (1994, 1999; advisory panel) suggests that the experience of violence results in a loss of voice and a loss of connection to others. In particular, she notes that connection to others is fundamental to girls' self-esteem. In looking outward for definition, in trying to please others, in being silenced, and in undergoing related life changes, girls can experience a "complete collapse on the inside." Friedman observes in addition, that "if you've been abused, one of the things that happens is your connections are severed – the trust, the connections with others." It is this severance of connection and loss of self that Friedman views as a fundamental cause of eating disorders.

Nine service providers told us that eating disorders are one way in which girls and women who have been abused seek to gain control over their lives. For example, one informant explained that

"The genius of oppression is that it denies us any control over our own lives except the power to destroy ourselves. Particularly, we are denied control over our own bodies."

- service provider and survivor of anorexia

"eating disorders are really about control and feeling powerless in a violent family." Indeed, research has demonstrated that sexual abuse history is linked to a lower level of perceived control (Waller, 1998). The service providers indicated that a lack of safety and security in the home may lead young women to search for an area of life that they can control. As one informant noted "the one thing they can control is their food intake." One front-line worker providing services to battered women noted that she frequently encounters "how food is regulated or denied or deprived, or other forms of using food as a weapon to control" and the internalization of this control by women.

The binge-purge cycle provides a means for "expressing of anger, relieving stress and tension, regaining a sense of self, establishing control, ensuring predictability and personal space, and 'cleansing' oneself of the abusive experience" (Everill & Waller, 1995b). For both the girl still living in the home and the woman who has left the parental home, food refusal may be one way for expressing hostility or punishment towards the abusing parent/caregiver and/or the parent who was unable to protect her (Brown, 1997; Lee & Lee, 1996; Williams, Wagner & Calam, 1992). The repeated violation of personal

body boundaries can lead to a sense of powerlessness in the woman (Everill & Waller, 1995a). She may feel that she has lost control of her body (Miller et al., 1993a) or that her body betrayed her by being weak, vulnerable, sexual, or small (Brown, 1997). "Both abuse and eating problems significantly disrupt a woman's ability to see her body as her own" (Thompson, 1994: 46-47). Levels of perceived control have been shown to be lower in women who report abuse and to be especially lower when attempts to regain control through disclosure have failed (Everill & Waller, 1995a). Ritualized eating behaviour is one way in which the survivor may impose a sense of control in her life and on her body (Brown, 1997). Whereas past psychodynamic theories have characterized this exertion of control as immaturity and teenage rebellion, more current models propose that the search for control is a creative and powerful act of resistance in an environment in which escape is impossible (Thompson, 1992, 1994).

Root (1996) has suggested that for groups who have been denied food through war or discrimination, eating symptomatology may manifest as food hoarding or compulsive eating. Prisoners who survived the last stage of starvation disease in Auschwitz-Birkenau, for instance, report a continued preoccupation with food (Ryn, 1990). Focus group discussions have suggested a relationship between African American eating patterns and the legacy of slavery (Airhihenbuwa et al., 1996). The generational effects of these experiences on eating should be considered as factors in assessment and treatment. The relationship to food may be more than just a healing process of the individual patient. The girl or woman's relationship to her community and her community's own healing processes around food and violence may be important to her recovery. One theorist in Prince George, B.C., for example, has argued that the restoration of the potlatch is central to Tsimshian identity, kinship, and heritage, as well as to recovery from the impact of colonialism (McDonald, 1995).

SELF-ESTEEM

Four service providers pointed to low self-esteem as a shared connection between eating disorders and a history of family violence. Three of these informants observed that abuse often leaves children with lower self-esteem, and eating disorders are one way of coping with these feelings. Indeed, eating disordered women who report unwanted sexual experience are more likely to see themselves as contaminated by the experience and to have lower sexual self-esteem (Waller et al., 1995). The shame that a woman feels about unwanted sexual experience has been connected to the development of anorexia (Schmidt, Tiller, Blanchard, Andrews & Treasure, 1997). Sexual abuse can result in the woman feeling disgusted with her own body, femaleness, and sexuality (Brown, 1997). Girls who are surviving or have survived childhood sexual abuse are more likely to see themselves as fat, ugly, and unworthy during adolescence (Schaaf & McCanne, 1994). Negative feelings are most directed toward those areas of the body associated with sexuality such as the breasts, stomach, and buttocks (Miller et al., 1993a) and can lead to dieting and purging in an effort to change these body parts (Thompson, 1992). Indeed, in research by Griffiths and McCabe (2000), adolescent girls scoring higher on body dissatisfaction showed higher levels of disordered eating behaviours and one of the most important predictors of body dissatisfaction was self-esteem.

One service provider noted that criticism of appearance can further undermine the self-confidence of girls and women. Thompson (1992: 556) has observed that “exposure to trauma did much more than distort the women’s visual image of themselves. These traumas often jeopardized their capacity to consider themselves as having bodies at all.” Thompson now uses the term “body consciousness” rather than “body image” to describe the woman’s ability to reside comfortably in her own body and to consider her body connected to herself. It is this body consciousness that is disrupted by abuse and which in turn linked to the development of eating disorders.

Bulimia has been linked to self-degradation among girls and women who are survivors of family violence (Everill & Waller, 1995b). Feelings of inadequacy, guilt, anxiety, inferiority, worthlessness, self-blame, and shame may provide the “psychic link” between abuse and the development of eating disorders (Brown, 1997; Hernandez, 1995). The guilt and self-blame that survivors experience is often further exploited by the abuser in an effort to maintain secrecy (Everill & Waller, 1995b).

THE SOCIOCULTURAL CONTEXT

The intersection of violence and eating disorders must be understood within the sociocultural context of women’s experiences. The diet and beauty industry sees gross earnings of approximately \$30 billion each year (Pike & Walsh, 1996). In an examination of 222 studies over the last 50 years, it was found that the number of women who have a poor body image compared to men has risen dramatically since the 1970s (DeAngelis, 1997). It has been argued that “comparing one’s own body to cultural ideals, and knowing one’s body will be subject to such comparisons by others, is fundamental to women’s experience” (Frederickson & Roberts, 1997: 192). In fact, eating practices and disordered eating in young and middle-aged women have been shown to be related to the extent to which they feel negatively about themselves when cultural body standards are not achieved (McKinley & Hyde, 1996).

Stereotypes regarding fatness have been repeatedly documented to begin early in childhood. Lawson (1980) found that 2nd, 4th, and 6th grade children had strong negative stereotypes about fat body figures and positive ones about average-sized figures. This trend worsened with increasing age. Similar responses were found in kindergarten through 4th grade by Brylinsky and Moore (1994). Research on adolescent girls found that fat models in photographs were evaluated less positively and less liked unless they were ascribed an “excuse” for their fatness (DeJong, 1980). In other words, there is an onus on the fat girl or woman to prove that her size is a medical problem and not an external symptom of a lazy, undisciplined, or disordered character. Indeed, Chrisler (1994) has observed that: “Image has become a central part of identity and is now used to convey invisible aspects of identity such as personality traits” (p. 226). Both fat and nonfat children have been found to generate more negative stereotypes of fat people (Counts & Jones, 1986). Degradation of fatness also exists among Canadian children and adolescents (LeBow, 1988; LeBow, Ness, Makarenko, & Lam, 1989).

Research has found that women's exposure to thinness-depicting media (magazines and television) predicts eating disorder symptomatology, a drive for thinness, body dissatisfaction, and a feeling of ineffectiveness (Harrison & Cantor, 1997). Young women are exposed to 10 times as many thinness-promoting advertisements as young men (Andersen & Holman, 1997). Media representations promote stereotypes that "women of colour are either fat and powerless (African American and Latina women); fat, bossy, and asexual; corrupt and/or evil (Asian/Pacific Americans, Island and African Americans); exotic (Asian American, mixed race); or hysterical and stupid. American Indian and Alaskan Native women are virtually nonexistent" (Root, 1990: 530).

By Grade 3, 40% of girls think that they should be thinner (Kraft, 1998). In Canada, more than 50% of girls under 18 years of age see themselves as too fat, even though 80% report a "normal" weight (McCreary Centre Society, 1993). Twice as many girls as boys see themselves as fat and as early as grades 3 to 6, twice as many girls as boys are dieting (Andersen & Holman, 1997). Of those who diet, 70 to 80% are dissatisfied with their bodies (Andersen & Holman, 1997). It is for this reason that some researchers have called for caution in speaking about the woman with anorexia or bulimia. Focusing predominantly on biomedical explanations, cognitive deficits and biases, maladaptive attitudes and beliefs, and physiological dysfunctions risks treating the woman solely as the source of the problem without also examining her social context (Malson & Ussher, 1996).

While the girl or woman receives messages about her appropriate body size and presentation, she is also exposed to portrayals of the violence that she may be experiencing in her life. Popular women's magazines, for example, predominantly characterize woman battering as a private problem and, specifically, the woman's problem (Berns, 1999). Similarly, reality-based television crime programmes tend to ascribe blame to the abused woman or represent her as uncooperative (Carmody, 1998). It is therefore not surprising that exposure to media aggression leads to more intense feelings of disempowerment for women (Pryor, Everett, & Ridener, 1999; Reid & Pinchilescu, 1995). Moreover, news reports about the rape and murder of women are most frequently written in the passive voice, which has been shown to result in a greater acceptance of violence among both women and men (Henley, Miller, & Beazley, 1995). The impact of these media representations is further compounded when projected into the woman's home where she may be surviving abuse. For the girl who is being sexually abused, she is confronted with a rising skepticism in the media reporting of child abuse (Benatar, 1995).

The girl or woman living with abuse is faced with a complex interaction of violence, verbal degradations, beauty standards, dieting and thinness norms, and media images that stigmatize her or disbelieve her abuse. In this context, her body becomes a shameful, sexualized, and "not-good-enough" body. It is blamed for the violence inflicted on it, as well as the way in which it grows and develops. The messages she receives from the media may be reinforcing the very messages she is being given by her abuser. Eating disorders may be one way in which the girl or woman seeks to gain control in relation to both her violated body and her discursive body.

ISSUES IN TREATING VIOLENCE AND EATING DISORDERS

Eight themes emerged from the responses by service providers to questions regarding their current treatment approaches and recommendations for treatment: integration of treatment; self-esteem; individualizing treatment; barriers of marginality; geographic isolation; safety; developmental treatment issues; and lack of funding and services.

INTEGRATION OF TREATMENT

Three of the service providers told us that the connection between family violence and eating disorders must be recognized and treated directly. An additional fourteen (11%) reported addressing this link as a central part of their approach. About 27% said that they address the connection if it arises. A further 5% provide an integrated treatment if the woman or girl herself prioritizes the connection as her therapeutic goal. Two programmes are developing information resources and treatment toward an integrated approach.

On the other hand, seven service providers commented that they do not alter their treatment approaches to either violence or to eating disorders if the two occur together. These providers indicated that abuse, for example, would be another issue to be addressed in healing self-esteem, but that it was not a central focus. Ten of the service providers reported approaching the treatment of abuse and eating disorders separately or sequentially. In some cases, the woman is transferred after the completion of one programme into another or is temporarily transferred for immediate help and then returned to the original programme. In other cases, she may first receive counselling for abuse and then receive counselling for eating disorders. Many women and girls who seek treatment for eating disorders and subsequently disclose abuse are transferred or referred. They are sent to other programmes, psychiatrists, abuse-related individual counselling, women's shelters, and/or the facilities of larger cities or provinces. In other cases, the advisory panel informed us, eating disorder programmes will not accept individuals who demonstrate co-morbidity, such as depression. This can further fragment care since, as one of our advisory committee members indicated: "It divvies the person up and sends their various symptoms to different clinics housed in the same individual." Yet, as another advisory committee member noted, this policy does not aptly reflect the circumstances of the woman: "What doesn't make sense is if a woman with an eating disorder and who is perhaps also experiencing violence in her intimate relationship, is not depressed."

Overall, there are relatively few resources that are established to directly assess, treat, and support girls and women who experience both abuse and eating disorders. Service providers indicated that they were eager for information, discussion, and resources addressing the links between abuse, anorexia, and bulimia. In the published literature, reviewers have emphasized that suggestions for an integrated treatment approach are sorely needed (Brown, 1997).

SELF-ESTEEM

Service providers and members of the advisory committee drew attention to the power relationship that exists between professionals and patients. One participant noted “going to the hospital is the most powerless experience a girl can have,” cautioning that this loss of power can risk heightening the girl’s need for her eating disorder. Another participant observed:

I’ve seen that things have been done wrong in hospitals where physicians and nurses have been in the position of control and power over very ill eating disordered people and have, in my considerable experience, frequently treated these women quite poorly.

Eleven service providers identified self-esteem as a core focus in treatment. As one informant explained, a programme for the woman or girl who has experienced both abuse and disordered eating must help build confidence, trust, and hope. It is important that the therapist-client relationship be warm and supportive and not recapitulate the controlling dynamics of past abusive relationships.

Service providers emphasized that clients need to be provided with new, positive coping skills for dealing with family violence and stress that can then replace the use of anorexia and bulimia. For example, the use of positive affirmation techniques and positive self-criticisms was suggested. A primary focus by the informants was on alternative ways in which the woman or girl could take control of her life or situation. One particularly critical point was made by two informants who noted the importance of identifying abusive relationships by naming the abuse and condemning the actions of the abuser. Re-framing an understanding of the violence away from self-blame is central to the re-visioning by the woman of her own self-worth and bodily integrity. In addition, women might be taught how to find resources for dealing with abuse and encouraged in assertiveness to overcome the fear of violence. Fourteen service providers addressed body image as a part of treatment and 21 provided the woman with education and awareness about violence, eating, nutrition, and/or body image.

INDIVIDUALIZING TREATMENT

Consistent with the research (Fichter, 1995), 10 of the service providers highlighted an approach to treatment which is individually based. Recovery was described as “a very individual process” and emphasis was placed on the girl or woman’s ability to prioritize the particular issues she would like addressed. In this way, each woman is invited to “make her own connection with her experiences.” Several informants noted that an integrated approach to treating abuse and eating disorders was used only when the client identified this as her goal. Some service providers encouraged women to take action in a way that was individually appropriate in order to aid healing. Suggestions for action included role playing, letter writing, and possible confrontation of the abuser. For some, the focus of counselling is on personal growth. Women are taught to “listen to their bodies” and to examine disordered eating as “a vehicle for learning about the self.”

BARRIERS OF MARGINALITY

According to the information we received from the service providers, available Canadian treatment and resources are offered predominantly in English only, with limited access to other languages. Fourteen of the service providers consulted offered services in either French only or in both French and English. Eight were able to provide help in languages other than French and English and 8 were able to access interpreters if required. One informant found the use of interpreters difficult when confidentiality or disclosure were critical issues. As was concluded by one service provider, those women who are not able to explore the issues of abuse and eating disorders in English "do not have much access and are generally restricted to treatment within their own communities."

While a variety of ethnic groups had been seen by service providers overall, many informed us that they served a rather homogeneous European-based clientele. Ten percent characterized their clients as "diverse." Some service providers specifically noted cultural barriers to receiving treatment, a reluctance to seek help by non-European women, and difficulty confiding in professionals who are unfamiliar with issues that these women face. Eurocentrism in the health care system, for example, has resulted in a lack of information about health issues affecting women of colour and has discouraged women from seeking help for fear of reinforcing racial stereotypes of themselves as "nurturing, well-nurtured, and overweight" (Bowen et al., 1991: 134; Dolan, 1991). Moreover, anorexia and bulimia have been mistakenly considered "the Golden Girl's Disease" (Root, 1990: 525), and women of colour have been assumed to be immune or "buffered by cultural differences" (DeAngelis, 1997; leGrange et al., 1998; Williamson, 1998). As a result, women of colour are often disbelieved, undiagnosed, misdiagnosed, and late diagnosed for eating disorders, leading to greater severity prior to diagnosis (Pike & Walsh, 1996; Thompson, 1992). This is especially problematic given that the severity upon presentation for treatment appears to be the best indicator of eating disorder prognosis (Casper & Jabine, 1996). Many women of colour have also been confronted with stereotypes of their communities as more prone to family violence, more patriarchal, and backwards or regressive in gender relations. As a result, women of colour may hesitate to disclose past or ongoing abuse to European service providers, for fear of exacerbating racial stereotypes, losing community, or betraying community solidarity (Flynn & Crawford, 1998).

Service providers also expressed concerns that they were not equipped to address the needs of Aboriginal women and girls. One informant noted that racism was a frequent part of the experiences of First Peoples who consulted her for eating disorders. The definition of "family violence" itself must be examined within a context of colonialism to include, for example, the history of residential schooling or apprehension from families and communities. "As adults or elders, individuals may feel that they do not have an avenue for voicing concerns or that what they experience is labelled as family violence, particularly in areas of emotional and mental abuse" (Frank, 1992: 4). It is important that service providers recognize that "family violence" may include institutional violence, violence imposed upon the family, and forms of systemic or oppressive violence within which the family must live.

Research has shown that lesbians also tend to avoid mainstream avenues to health care and turn more often to alternative health because of past discriminatory experiences (Simkin, 1991, 1998). Lesbians who do seek help may avoid identifying their sexuality for fear of homophobia (Ministry of Health and Ministry Responsible for Seniors, 1995). Most lesbians report wanting to disclose their sexuality to family physicians, but need to feel safe and secure to do so (Geddes, 1994 in Simkin, 1998). In addition, lesbians who report current or past battering by a partner or sexual assault by a woman need to be believed. Practitioners require an understanding of the issues specific to lesbian battering that differ from heterosexual women's experiences – lack of resources and safe shelter, community loyalties, lack of anonymity within the community, disbelief by authorities and by community, and internalized lesbophobia. The disbelief of violence within lesbian relationships may be compounded by assumptions that lesbians have a “protective buffer” against societal standards of female beauty and against disordered eating (DeAngelis, 1997). Lesbians are not immune to the cultural standards in which they live (Heffernan, 1996). The belief that they are not generally at risk for anorexia, bulimia, battering, or woman-on-woman sexual assault can lead to under-recognition of problems and to alienation of the woman within treatment.

One service provider noted inaccessibility and lack of informed services for women with disabilities/chronic illnesses. Yet, women and girls with disabilities face higher rates of physical abuse, sexual abuse, and battering (Health Canada, 1996; National Clearinghouse on Family Violence, 1997b). Moreover, hearing impaired women do not differ from hearing women in attitudes toward food or dieting behaviour (Fletcher, 1993). Women with disabilities/chronic illnesses are already stigmatized as the “abnormal body” (Davis, 1995) and risk further pathologizing in disclosing disordered eating. Medical approaches to anorexia and bulimia in the absence of therapeutic support may further impose procedures on an already over-medicalized body, thereby discouraging some women with disabilities/chronic illnesses from seeking treatment. Women with disabilities/chronic illnesses who present with eating disorders need to be asked about the presence and role of past or current abuse. In addition, women with disabilities/chronic illnesses who seek help for abuse and display signs of disordered eating should not be assumed to be displaying a simple “medical complication” and have their symptoms dismissed.

It must be recognized by clinicians and researchers that eating disorders affect heterosexual and bisexual women and girls and lesbians of all racial, ethnic, and class backgrounds. Primary care providers, social service providers and other professionals must challenge the myth that disordered eating impacts only or mostly white, middle-class and/or heterosexual girls and women. Both research and treatment projects need to be proactive in including girls and women of colour (Root, 1990), lesbians, and women/girls living in poverty. Clinicians must take the initiative to become culturally literate and the time to build community relationships (Root, 1990). Culturally-specific treatment will consider the impact of historical and current racism, immigration experiences, institutionalization, and colonialism on communities and individuals. In particular, the impact of food deprivation across generations and for those currently living in poverty

must be further researched and be factored into individual treatment where applicable. Moreover, service providers must be sensitive to both disability and to abuse. Physically accessible services are necessary, as well as sensitivity to the needs of girls and women with hearing, visual, cognitive, and developmental disabilities.

"The core psychological themes reflected in disordered eating are the pursuit of identity, power, specialness, validation, self-esteem, and respect - themes significant in the lives of all oppressed persons."

- Root, 1990: 526

GEOGRAPHIC ISOLATION

Women living in isolated, rural, or small communities who are experiencing violence face unique barriers to accessing treatment: a lack of anonymity within their home areas, the threat of firearms, lack of public or personal transportation, lack of protective services such as police within their areas, and an inability to physically leave the area or an incident of battering (Jiwani with Moore & Kachuk, 1998). Such factors may compound the sense of powerlessness and lack of control that these women feel. Control over eating may become one way in which to manifest a sense of self. There may be few services for girls and women seeking help for abuse and eating disorders. For example, some regions reported no publicly-funded programmes, or limited access to psychologists and nutritionists. Women are referred to services in the closest large city, which can be an hour or more away. Small urban centres reported that women and girls requiring more intensive treatment or hospitalization must be moved to larger cities. For smaller provinces and for northern areas, women and girls must often be sent out of province, away from community support and family, to receive appropriate help. Service providers noted that this is especially true in acquiring services that address the connections between family violence and disordered eating. Counsellors are also more likely to have to travel in order to meet the needs of the community and surrounding area.

SAFETY

A few service providers reported that they do not address violence for women and girls with anorexia and bulimia unless their client is in danger from her abuser. Those service providers who used an integrated approach, however, stressed that the individual's safety is a prime concern during therapy and in considering programmatic changes. Suicide checks and assessments of risk to ensure the safety of the girl or woman from violence were recommended. One informant noted that, upon disclosure of abuse history in women with eating disorders, he avoided doing physical examinations beyond routine requirements in order to avoid further traumatization. Similarly, another service provider noted that a history of abuse affected her approach to body work in treatment. Safety issues are emphasized more "in an attempt to have clients feel safe inside their bodies."

DEVELOPMENTAL TREATMENT ISSUES

Approximately 5% of all cases of anorexia develop among children younger than 12 years old (Nielsen, Lausch & Thomsen, 1997). The child's accessibility to treatment is a particularly pertinent issue when she is presenting with both disordered eating and a history of family violence. As one service provider indicated, abuse must be reported, and the presence of abuse determines whether state child welfare agencies will become involved. Often parents will try to withdraw children from treatment as a consequence. Similarly, another informant explained that some parents or guardians may not cooperate with treatment because one parent/caregiver is the perpetrator of abuse or because divorce affects relationship dynamics. But researchers have noted that many children who are abused show difficulties with eating in the short term and early intervention may be critical in preventing the development of eating disorders (Williams et al., 1992).

The Canadian Pediatric Society (1998) has highlighted some of the barriers to effective care for adolescent girls. Provincial health care plans often place limits on access to private care resources such as mental health or nutrition counselling. In addition, as the adolescent ages, she may no longer be eligible for coverage under provincial medical insurance rules. She therefore finds her treatment ending at a time when she is faced with such developmental challenges as leaving home, unemployment, or temporary employment. Moreover, she may be forced into transition from pediatric to adult care by the age limit policies of some treatment programmes. One service provider noted a lack of services for those individuals who are in transition from youth to adult programmes. For some, such disruption in care may hinder or endanger recovery. Youth health centres were suggested as important locations for early identification and prevention programmes (advisory panel). Important to this provision, however, is the need for confidentiality and accessibility, as many youth may be unwilling to disclose violence or eating disorders to family or meet with service providers if it requires transportation arrangements (advisory panel).

For some older women, anorexia may replace previous forms of self-harm, such as wrist cutting (Nicholson & Ballance, 1998). For some, it may be a relapse to earlier eating disorders. "No former sufferer should be considered immune from relapse at any age,

especially following a bereavement or other significant loss” (Cosford & Arnold, 1992: 497). Clinicians should be alert to previous histories of eating disorders, depression, battering, sexual abuse, and post-traumatic stress. Some older women who enter treatment for eating disorders may be reluctant to discuss childhood sexual abuse and sexuality issues (Beck et al., 1996). Recurrence or late onset of eating disorders may be triggered by the death of somebody close (e.g., a spouse), institutionalization, loss of independent living, or moving in with family. For older women, limited financial resources may determine poor diet and inability to access services. It is important that services addressing the intersections of violence and disordered eating be available to aging women. Such services must include consistent and appropriate screening.

LACK OF FUNDING AND SERVICES

A consistent problem identified by service providers was the impact of increasing funding cutbacks and of lack of funding. Treatment facilities are facing numerous difficulties in continuing to meet the needs of individuals who have experienced abuse and who are at risk from anorexia and bulimia. Service providers across Canada told us that they were struggling to survive staff shortages, decreased programme lengths, free and frequent staff overtime, expanded geographic areas to serve, re-allocation of treatment time to fundraising, and increasing reliance on volunteers. One service provider was forced to work from home in order to reduce costs. Service providers told us that, for many services, there is pressure to avoid treating the women who are most seriously ill, and thus most at risk, because of higher treatment costs. While some services are facing urgent financial crises and are in danger of closing, others have already shut down all services or are on hold. The ability to be inclusive of marginalized women and girls has been compromised and, in many cases, those individuals must pay for private treatment at \$100 per hour for individuals and \$2400 for a 4-month group programme. Some facilities that were previously addressing both violence against women and eating disorders have now restricted their services solely to the treatment of eating disorders, due to funding cuts. A residential programme reported having only partial funding that covered neither the cost of residency nor of food. Piecemeal and inconsistent funding disrupts the effectiveness of programmes and means that staff must concentrate their limited time on grant writing and fundraising. As a result, the ability to focus on treatment itself is diminished. Waiting lists into programmes range from 1 week to 6 months for emergency cases and can be even longer for non-emergencies.

Service providers reported that as a result of budgetary cutbacks, programmes and services are being downloaded onto community groups, most of which lack sufficient resources to respond. Community-based groups are often closing or being reduced to referral services. Hospital services are increasingly being limited to those patients who are at high risk of dying. As a result, forced weight gain is too frequently the exclusive focus of medical treatment, without the necessary after-care or psychotherapeutic intervention. Underlying issues of family violence, abuse, powerlessness, family dynamics, and control are left unaddressed. It is crucial to the eating disordered girl or woman who has experienced abuse that she be approached as a *whole* person in order for treatment outcomes to be lasting and effective.

KEY COMPONENTS IDENTIFIED FOR THE TREATMENT OF VIOLENCE AND EATING DISORDERS

Approximately two-thirds of the service providers reported a connection between abuse and disordered eating. Almost half of the total sample characterized this link as strong. Of these, all except one attempted to use an integrated approach to treatment, either as a central component or as a priority identified by the client. Many of our participants, however, emphasized the need for more information and resources. In addition, they offered suggestions for treatment.

Based on a review of research and on the input of Canadian service providers, twelve therapeutic concerns across four stages of treatment have been identified as necessary to an integrated treatment approach to disordered eating and violence against women and girls.

| INTEGRATED TREATMENT | |
|-----------------------------|-----------------------------|
| STAGE OF TREATMENT | THERAPEUTIC CONCERNS |
| PRESENTATION | ACCESSIBILITY |
| | ASSESSMENT |
| IMMEDIATE NEEDS | SAFETY |
| | MEDICAL STABILITY |
| THERAPEUTIC SUPPORT | VALIDATION |
| | SELF-WORTH |
| | RE-FRAMING |
| | COPING SKILLS |
| | CONTROL & EMPOWERMENT |
| | RELATIONSHIPS & NETWORKS |
| | EDUCATION & AWARENESS |
| CONTINUITY | FOLLOW-UP |

PRESENTATION

1. Accessibility

Girls and women must have access to adequate services, appropriate services, and to already existing services. The provision of access means that:

- issues pertinent to marginalized women are understood and addressed in treatment
- facilities are physically accessible to and specialized for girls and women with disabilities/chronic illnesses
- both abuse and eating disorders are properly identified across race, class, language, ancestry, and developmental lines
- services are affordable and sufficiently funded
- sufficient community-based programmes exist within small and rural areas.

Maria Root (1990) has called on health professionals to take the initiative to be culturally literate, to actively participate in organizations that are multiracial and multicultural, and to be aware of cultural understandings of causation, help-seeking, and cure. She has highlighted the importance of building relationships within various communities and learning from key community members. Practitioners have not, for example, expected, looked for, or treated anorexia or bulimia among homeless people, despite higher rates of both a history of abuse and of eating disorders in this population (Gard & Freeman, 1996).

Members of the advisory panel emphasized that transition houses are important sites for intervention, noting that: “it’s a residential setting so there are lots of ways to make things accessible whether it’s education about body image, body size, body consciousness, or just the food that’s available. For lots of women who go to transition houses, it’s the first time that they can actually eat freely and eat what they want.”

2. Assessment

Several recommendations emerged for assessment:

- Service providers who are working with girls and women experiencing sexual, physical, and emotional family or institutional (e.g., foster homes, residential schools) violence should inquire carefully about eating patterns (Williams et al., 1992).
- Girls and women presenting with eating disorders should be asked directly about past and present experiences of violence, emotional abuse, and abusive interactions surrounding food. Either interviewing or a combined use of admission forms and interviewing may be used, structured with overlapping, open-ended questions.

- The presence of eating disorders, childhood abuse, and partner abuse should be assessed across age and sexual identity (e.g., battering in lesbian and/or elder couples, adolescent battering, the impact of child abuse for older women).
- The messages that the girl or woman has received about her body from family should also be explored, including the impact of family changes in class positioning, “coming out,” dieting, and cultural assimilation.
- Assessors should also check for possible accompanying or co-morbid risks, such as depression, post-traumatic stress disorder, self-harm, and suicidal ideation or attempts.

IMMEDIATE NEEDS

3. Safety

The girl or woman who is living with abuse should have a safety plan. This includes protection from reprisals, punishment, or being withdrawn from treatment by the family or abuser. The abused adult should be assured consistent support and appropriate safety measures if she makes the decision to confront, escape, or report the violence. Safety also includes resources for meeting basic needs. Individuals who are homeless or living in poverty will need appropriate and sufficient food, food storage facilities, and shelter to facilitate recovery. Women and girls with disabilities will require accessibility to both services and to safe houses/transition houses as well as attentiveness to higher levels of poverty. Joan Meister (1990), of the DisAbled Women’s Network (DAWN) has noted that: “Often the only cushion between a disabled woman and poverty on social assistance or poverty on minimum wage is the financial support of the family. Yet dependence on family is a demeaning and even dangerous place for many women with disabilities. ... Battering, incest, and other forms of abuse are endemic and unrecognized by the support services which are largely inaccessible” (p. 42).

4. Medical Stability

In conjunction with therapeutic support, medical stabilization of the body may be necessary. This includes access to medical treatment, monitoring, advice, and emergency hospitalization. In addition, support for addiction recovery may be required. Attention to the woman’s health needs should occur within a supportive, caring atmosphere that encourages her sense of wholeness or “body consciousness” (Thompson, 1992, 1994). Health practitioners need to be particularly aware of possible abuse histories among eating disordered women and be respectful in approaching the woman’s body. Body boundaries and issues of consent must be carefully considered in providing medical treatment.

THERAPEUTIC SUPPORT

5. Validation

Service providers must be aware of their own reactions to revelations of abuse by women or girls with eating disorders. Responses should be appropriately warm and empathetic. Expressions of horror, shock, disbelief, disgust, or complete neutrality may exacerbate shame, self-doubt, and self-blame (Everill & Waller, 1995b). It is important that the strength and survival of the woman be acknowledged, that the abuse be named, and that the woman's experiences be believed.

6. Self-worth

Counselling should be aimed at the restoration or promotion of a sense of the self as worthy, capable, and trustworthy. Self-affirmations, a more holistic "body consciousness," and more constructive forms of self-criticism should be developed to displace forms of self-denigration or internalized messages of oppression. One service provider suggested that the girl or woman be actively engaged in prioritizing issues and establish individualized goals within treatment.

7. Re-framing

The woman's beliefs about the abuse, about herself, and about her body should be openly discussed. In particular, self-blame should be identified and re-framed through the clear condemnation of the abuser's actions. Reactions to previous attempts she has made at disclosing should be discussed and inappropriate responses by caregivers, professionals, or family should be acknowledged.

8. Coping Skills

A number of service providers suggested teaching coping skills during counselling. For example, discussion might address alternative ways to reassert personal control, methods of establishing and enforcing new body boundaries, strategies for seeking resources or help when needed, the power to recognize and name abuse, and the creation of a political or social voice.

9. Control and Empowerment

It is critical to support the girl or woman in gaining the sense of control that she may be seeking. This might develop through re-defining her role, rights, and status within the family or, alternatively, helping her to leave the family. It may come through establishing a sense of herself as a strong individual. In addition, one service provider has observed that some women find "calling their abuser to account an important step toward healing." In particular, this informant noted that some women have found phoning their abusers, writing them letters, confronting them in person, or lodging a complaint with police a "powerful form of self-validation." In research by Roush (1999), most of the incest survivors who confronted their abusers characterized the action as empowering and important to their healing. These same women also noted a mixture of family reactions from

improvement to terminated relationships and a mixture of personal reactions, including shock, fear, anger, and regret. As noted by one advisory panel member and by Cameron (1994), the woman's safety must be paramount. Moreover, the advisory panel noted that eating disorder symptoms may flare up in response to the increased risks in which the woman or girl sees herself engaging during the recovery process.

10. Relationships and Networks

The establishment of healthy relationships, support networks, and community membership/relations should be a part of the therapeutic goal. This may be variously accomplished through:

- family therapy
- support groups
- building of relationship skills
- helping the woman to define her own goals for a healthy relationship
- establishing a network of community advocates and resources that the girl or woman can contact or join
- ensuring that the therapist-client relationship is supportive and does not re-create the power dynamics that the woman has experienced during abuse
- supporting the woman in her relationship to her own racial, ethnic, aboriginal, sexuality, disability/chronic illness, reserve, peer, or rural community and understanding the role that she wishes her community to play in her healing.

11. Education and Awareness

Service providers recommended that the recovery process include consciousness raising about media, family, peer, and other sociocultural influences on body image and in issues of violence. The provision of libraries, videos, workshops, and peer discussion groups can challenge the stigmatization of fatness, disordered eating, sexuality, addiction of self or family, sexual abuse, and physical abuse that creates shame and silence.

CONTINUITY

12. Follow-up

It is critical that the recovering girl or woman be able to retain a sense of connection, receive counselling subsequent to medical stabilization, and receive follow-up checks for health complications or relapse. Provisions might include check-in calls, letters, or cards, drop-in access, available telephone support, continued access to support groups, gradual release, and the opportunity to become involved in educational or peer counselling programmes. Funding cutbacks, however, have made treatment programmes themselves difficult to sustain and have eliminated the necessary follow-up from many programmes.

CONCLUSION

The literature and the data obtained from interviews with 123 service providers identified a relationship between eating disorders and violence against women and girls. This finding was affirmed by feedback from our national advisory committee. However, both the literature and the interview data caution us that the connection is not a simplistic or an inevitable one. Rather, the particular experiences of the individual girl or woman will shape the likelihood, nature, and personal meaning of eating disorders in her life. For some women, eating disorders may represent resistance against past and further violation of the body. For others, they may represent a way to destroy the body and purge feelings of shame and guilt. Exploring the specific importance of anorexia or bulimia as a response to abuse for the individual is central to recovery. It is a focus on the unique needs of the girl or woman in assessment and treatment that characterizes an *integrated approach*. Specific components of an integrated approach suggested here include an emphasis on accessibility; assessment; safety; medical stability; validation; self-worth; re-framing; coping skills; control and empowerment; relationships and networks; education and awareness; and adequate follow-up.

Although research is increasingly examining the manner in which abuse may contribute to the development of disordered eating, there has been relatively little focus on resiliency factors within this relationship. A direction for future research posed by the advisory committee of this project is the investigation of the variables or circumstances that “inoculate people or provide them with supports and coping strategies” in averting the development of eating disorders.

WORKS CITED

- Airhihenbuwa, C.O., S. Kumanyika, T.D. Agurs, A. Lowe, D. Saunders & C. Morssink. "Cultural Aspects of African American Eating Patterns." *Ethnicity and Health*, 1 (1996): 245-260.
- Alliance of Five Research Centres on Violence. *Violence Prevention and the Girl Child, Final Report*. Vancouver, BC: Alliance of Five Research Centres on Violence, 1999.
- American Psychiatric Association. *Diagnostic and Statistical Manual of Mental Disorders: DSM-IV*. Washington, DC: APA, 1994.
- Andersen, A.E. & J.E. Holman. "Males with Eating Disorders: Challenges for Treatment and Research." *Psychopharmacology Bulletin*, 33 (1997): 391-397.
- Andrews, B. "Bodily Shame in Relation to Abuse in Childhood and Bulimia: A Preliminary Investigation." *British Journal of Clinical Psychology*, 36 (1997): 41-49.
- Anorexia and Bulimia Nervosa Foundation of Victoria, Inc. June 2000.
Available online:
http://www.nevdgp.org.au/geninf/std_misc/ANOREXIA_AND_BULIMIA_NERVOSA.htm
- Atwood, N.C. "Gender Bias in Families and Its Clinical Implications for Women." *Social Work*, 46 (2001): 23-36.
- Baldo, T.D., W. Bostwick & A.J. Baldo. "Intrafamilial Assaults, Disturbed Eating Behaviours, and Further Victimization." *Psychological Reports*, 79 (1996): 1057-1058.
- Baldo, T.D., W. Bostwick, S.D. Wallace & M.S. O'Halloran. "Effects of Intrafamilial Sexual Assault on Eating Behaviours." *Psychological Reports*, 79 (1996): 531-536.
- Battered Women's Support Services. *Violence Against Women in Relationships in British Columbia: A Fact Sheet*. Vancouver: BWSS, 1997.
- BC Institute Against Family Violence. October 2001. Available online:
<http://www.bcifv.org/about/overview/3.html>
- BC Ministry of Health and Ministry Responsible for Seniors. *A Report on the Health of British Columbians: Provincial Health Officer's Annual Report. Feature Report: Women's Health*. Victoria: Office of the Provincial Health Officer, 1995.
- Beatus, J. "Body Image, Self-Concept and Factors which Influence the Adjustment to Disability in Spinal Cord Injured Persons." *Dissertation Abstracts International*, 57, 7-B (1997): 4334.
- Beck, D., R. Casper & A. Andersen. "Truly Late Onset of Eating Disorders: A Study of 11 Cases averaging 60 Years of Age at Presentation." *International Journal of Eating Disorders*, 20 (1996): 389-395.

- Beckman, K.A. & G.L. Burns. "Relation of Sexual Abuse and Bulimia in College Women." *International Journal of Eating Disorders*, 9 (1990): 487-492.
- Ben-Tovim, D.I. & M.K. Walker. "Body Image, Disfigurement, and Disability." *Journal of Psychosomatic Research*, 39 (1995): 283-291.
- Benatar, M. "Running Away from Sexual Abuse: Denial Revisited." *Families in Society*, 76, 5 (1995): 315-320.
- Benson, C. "The War on Women." In *Social Problems in Canada Reader*, ed. E.D. Nelson & A. Fleras, 381-392. Scarborough, ON: Prentice Hall Canada, 1995.
- Berns, N. "'My Problem and How I Solved It': Domestic Violence in Women's Magazines." *Sociological Quarterly*, 40, 1 (1999): 85-108.
- Bowen, D.J., N. Tmoyasu & A. Cauce. "The Triple Threat: A Discussion of Gender, Class, and Race Differences in Weight." *Women and Health*, 17, 4 (1991): 124-143.
- Brown, L. "Child Physical and Sexual Abuse and Eating Disorders: A Review of the Links and Personal Comments on the Treatment Process." *Australian and New Zealand Journal of Psychiatry*, 31 (1997): 194-199.
- Bryant-Waugh, R. & B. Lask. "Anorexia Nervosa in a Group of Asian Children Living in Britain." *British Journal of Psychiatry*, 158 (1991): 229-233.
- Brylinsky, J.A. & J.C. Moore. "The Identification of Body Build Stereotypes in Young Children." *Journal of Research in Personality*, 28, 2 (1994): 170-181.
- Cachelin, F.M. & B. Maher. "Is Amenorrhea a Critical Criterion for Anorexia Nervosa." *Journal of Psychosomatic Research*, 44 (1998): 435-440.
- Cameron, C. "Women Survivors Confronting Their Abusers: Issues, Decisions, and Outcomes." *Journal of Child Sexual Abuse*, 3, 1 (1994): 7-35.
- Canadian Centre for Justice Statistics. *Juristat Service Bulletin*, 14, 8 (1994): 10.
- Canadian Pediatric Society. "Eating Disorders in Adolescents: Principles of Diagnosis and Treatment." *Paediatrics and Child Health*, 3 (1998): 189-192. Available online: <http://www.cps.ca/english/statements/AM/am96-04.htm>
- Canadian Research Institute for the Advancement of Women. Available online: <http://criaw-icref.ca/>
- Carmody, D.C. "Mixed Messages: Images of Domestic Violence on 'Reality' Television." In *Entertaining Crime: Television Reality Programs*, ed. M. Fishman & G. Cavender, 159-174. New York: Aldine de Gruyter, 1998.
- Casper, R.C. & L.N. Jabine. "An Eight-Year Follow-Up: Outcome from Adolescent Compared to Adult Onset Anorexia Nervosa." *Journal of Youth and Adolescence*, 25 (1996): 499-517.
- Chrisjohn, R.D. & S.L. Young, with contributions by M. Maraun. *The Circle Game: Shadows and Substance in the Indian Residential School Experience in Canada*. Penticton, BC: Theytus Books Ltd., 1997.

- Chrisler, J.C. "Out of Control and Eating Disordered." In *Feminist Perspectives on Addictions*, ed. Nan Van Den Bergh, 139-149. New York: Springer Publishing, 1991.
- Clinique St-Amour: Le Centre de Traitement de l'Anorexie Nerveuse, la Boulimie et Autres Troubles de l'Alimentation. 2001. Available online: <http://www.cliniquestamour.com/>
- Connors, M.E. & W. Morse "Sexual Abuse and Eating Disorders: A Review." *International Journal of Eating Disorders*, 13, 1 (1993): 1-11.
- Cosford, P. & E. Arnold "Eating Disorders in Later Life: A Review." *International Journal of Geriatric Psychiatry*, 7 (1992): 491-498.
- Counts, C.R. & C. Jones. "The Perception of Obesity by Normal-Weight versus Obese School-Age Children." *Child Psychiatry and Human Development*, 17, 2 (1986): 113-120.
- Crago, M., C. Shisslak & L.S. Estes. "Eating Disturbances among American Minority Groups: A Review." *International Journal of Eating Disorders*, 19 (1996): 239-248.
- Crandall, C.S. & M. Biernat. "The Ideology of Anti-Fat Attitudes." *Journal of Applied Social Psychology*, 20, 3 (Part 2) (1990): 227-243.
- Daniels, J. "The Diagnosis of Eating Disorders in Women of Color." October 2001. Available online: http://www.something-fishy.org/cultural/roles_jd.php
- Danielson, K.K., T.E. Moffitt, A. Caspi & P.A. Silva. "Comorbidity between Abuse of an Adult and *DSM-III-R* Mental Disorders: Evidence from an Epidemiological Study." *American Journal of Psychiatry*, 155, 1 (1998): 131-133.
- Dansky, B., T.D. Brewerton, D.G. Kilpatrick & P.M. O'Neil. "The National Women's Study: Relationship of Victimization and Posttraumatic Stress Disorder to Bulimia Nervosa." *International Journal of Eating Disorders*, 21, 3 (1997): 213-228.
- Davenport, C., K. Browne & R. Palmer "Opinions on the Traumatizing Effects of Child Sexual Abuse: Evidence for Consensus." *Child Abuse and Neglect*, 18, 9 (1994): 725-738.
- Davis, C. & J. Yager "Transcultural Aspects of Eating Disorders: A Critical Literature Review." *Culture, Medicine, and Psychiatry*, 16 (1992): 377-394.
- Davis, L.J. *Enforcing Normalcy: Disability, Deafness, and the Body*. London: Verso, 1995.
- Day, D. *Young Women in Nova Scotia Survey*. Antigonish, NS: Cumberland County Transition House Association, 1990.
- DeAngelis, T. "Body-Image Problems affect All Groups." *APA Monitor* 1997. Available online: <http://www.apa.org/monitor/>

- DeGroot, J.M., S.R. Kennedy & G. McVey "Correlates of Sexual Abuse in Women with Anorexia Nervosa and Bulimia Nervosa." *Canadian Journal of Psychiatry*, 37 (1992): 516-518.
- DeJong, W. "The Stigma of Obesity: The Consequences of Naïve Assumptions Concerning the Causes of Physical Deviance." *Journal of Health and Social Behaviour*, 21, 1 (1980): 75-87.
- Deschamps, G., ed. *2-Spirited People of the 1st Nations: We are Part of a Tradition*. Toronto: 2-Spirit People of the 1st Nations, 1998.
- Deschamps, G. & S. Wahsquonaikeshik. "Two Spiritedness." In *2-Spirited People of the 1st Nations: We are Part of a Tradition*, ed. G. Deschamps, 10-11. Toronto: 2-Spirit People of the 1st Nations, 1998.
- DisAbled Women's Network (DAWN) Canada. *Beating the Odds: Violence Against Women with Disabilities*. Position Paper #2. Toronto: DAWN, 1989.
- Dolan, B. "Cross-Cultural Aspects of Anorexia Nervosa and Bulimia: A Review." *International Journal of Eating Disorders*, 10, 1 (1991): 67-78.
- Duffy, A. "The Feminist Challenge: Knowing and Ending the Violence." In *Feminist Issues: Race, Class, and Sexuality*, ed. N. Mandell, 132-159. Scarborough, ON: Prentice Hall Allyn and Bacon Canada, 1998.
- Dumont-Smith, C. & P. Sioui Labelle. "Dragging Wife Abuse Out of the Closet." Cited in Health Canada, *Breaking the Links between Poverty and Violence Against Women*, 1996, p. 41.
- Education Wife Assault. "The Impact of Cuts on Abused Women and Shelters for Women." *Education Wife Assault Newsletter*, 7 (1996).
- Everill, J. & G. Waller "Disclosure of Sexual Abuse and Psychological Adjustment in Female Undergraduates." *Child Abuse and Neglect*, 19, 1 (1995a): 93-100.
- _____. "Reported Sexual Abuse and Eating Psychopathology: A Review of the Evidence for a Causal Link." *International Journal of Eating Disorders*, 18, 1 (1995b): 1-11.
- Fichter, M.M. "Inpatient Treatment of Anorexia Nervosa." In *Eating Disorders and Obesity: A Comprehensive Handbook*, ed. K.D. Brownell & C.G. Fairburn, 336-343. New York: Guilford Press, 1995.
- Field, A.E., G.A. Colditz & K.E. Peterson. "Racial Differences in Bulimic Behaviours among High School Females 1-7." *Annals of the New York Academy of Science*, 817 (1997): 359-360.
- Finn, S., M. Hartman, G. Leon & L. Lawson. "Eating Disorders and Sexual Abuse: Lack of Confirmation for a Clinical Hypothesis." *International Journal of Eating Disorders*, 5 (1986): 1051-1060.

- Fletcher, M.L. "A Pilot Study of the Eating/dieting Behaviours and Attitudes of College-Aged Women Who are Hearing Impaired vs. Their Hearing Peers." *Journal of the American Deafness and Rehabilitation Association*, 27 (1993).
- Flynn, K. & C. Crawford "Committing 'Race Treason': Battered Women and Mandatory Arrest in Toronto's Caribbean Community." In *Unsettling Truths*, ed. K.D. Bonnycastle & G. S. Rigakos, 93-102. Vancouver: Collective Press, 1998.
- Ford, K.A. "Bulimia in an Egyptian Student: A Case Study." *International Journal of Eating Disorders*, 11, 4 (1992): 407-411.
- Fournier, S. & E. Crey. *Stolen from Our Embrace*. Vancouver: Douglas and McIntyre, 1997.
- Frank, S. *Family Violence in Aboriginal Communities: A First Nations Report*. Victoria: Ministry of Women's Equality, March, 1992.
- Frederickson, B.L. & T. Roberts. "Objectification Theory: Toward Understanding Women's Lived Experiences and Mental Health Risks." *Psychology of Women Quarterly*, 21 (1997): 173-206.
- Freeman, C.P. & M.C. Gard. *Eating Disorders in a Young Homeless Population*. Manuscript cited in M.C. Gard & C.P. Freeman, "The Dismantling of a Myth: A Review of Eating Disorders and Socioeconomic Status." *International Journal of Eating Disorders*, 20, 1 (1996): 1-12.
- French, S.A., M. Story, G. Remafedi, M.D. Resnick & R.W. Blum. "Sexual Orientation and Prevalence of Body Dissatisfaction and Eating Disordered Behaviours: A Population-Based Study of Adolescents." *International Journal of Eating Disorders*, 19 (1996): 119-126.
- Friedman, S.S. *Girls in the 90s: Facilitator's Manual*. Vancouver: Salal Books, 1994.
- _____. *Just for Girls: Facilitator's Manual*. Vancouver: Salal Books, 1999.
- Friedrich, W.N., A.J. Urquiza & R.L. Beilke. "Behaviour Problems in Sexually Abused Young Children." *Journal of Pediatric Psychology*, 11 (1986): 47-57.
- Gadd, J. "More Boys Physically Abused than Girls." *Globe & Mail*, July 9, 1997: A1, A6.
- Gagnon, L. "Despite Image, Most Anorexics are 45 or Older." *The Medical Post*, October 8, 1996. Available online: <http://www.mentalhealth.com/mag1/p5m-et01.html>
- Gard, M.C. & C.P. Freeman. "The Dismantling of a Myth: A Review of Eating Disorders and Socioeconomic Status." *International Journal of Eating Disorders*, 20, 1 (1996): 1-12.
- Garfinkel, P.E., E. Lin, P. Goering, C. Spegg, D.S. Goldbloom, S. Kennedy, A.S. Kaplan & D.B. Woodside. "Bulimia Nervosa in a Canadian Community Sample: Prevalence and Comparison of Subgroups." *American Journal of Psychiatry*, 152, 7 (1995): 1052-1058.

- _____. "Purging and Nonpurging forms of Bulimia Nervosa in a Community Sample." *International Journal of Eating Disorders*, 20, 1 (1996a): 231-238.
- _____. "Should Amenorrhea be Necessary for the Diagnosis of Anorexia Nervosa?" *British Journal of Psychiatry*, 168 (1996b): 500-506.
- Geddes, V.A. "Lesbian Expectations and Experiences with Family Doctors. How Much Does the Physician's Sex Matter to Lesbians?" *Canadian Family Physician*, 40 (1994): 908-920.
- Goldbloom, D.S. "The Early Canadian History of Anorexia Nervosa." *Canadian Journal of Psychiatry*, 42, 2 (1997): 163-167.
- Goldbloom, D.S., M. Olmsted, R. Davis, J. Clewes, M. Heinmaa, W. Rockert & B. Shaw. "A Randomized Controlled Trial of Fluoxetine and Cognitive Behavioural Therapy for Bulimia Nervosa: Short-Term Outcome." *Behavioural Research and Therapy*, 35 (1997): 803-811.
- Gowers, S.G., C.D. North, V. Byram & A.B. Weaver. "Life Events Precipitants of Adolescent Anorexia Nervosa." *Journal of Child Psychology and Psychiatry*, 37 (1996): 469-477.
- Grandmaison, A. *Protection des personnes âgées: Étude exploratoire de la violence à l'égard de la clientèle des personnes âgées du CSSMM*. Montréal: Centre de services sociaux du Montréal Métropolitain, Direction des services professionnels. Cited in Health Canada, *Breaking the Links Between Poverty and Violence Against Women*, 1996, p. 33.
- Griffiths, J.A. & M.P. McCabe. "The Influence of Significant Others on Disordered Eating and Body Dissatisfaction among Early Adolescent Girls." *European Eating Disorders Review*, 8 (2000): 301-314.
- Groupe d'Etude Francais sur l'Anorexie et la Boulimie. 2001.
Available online: <http://perso.club-internet.fr/dorigram/>
- Harris, S.M. "Racial Differences in Predictors of College Women's Body Image Attitudes." *Women and Health*, 21, 4 (1994): 89-104.
- Harrison, K. & J. Cantor "The Relationship between Media Consumption and Eating Disorders." *Journal of Communication*, 47 (1997): 40-67.
- Health Canada. *Breaking the Links between Poverty and Violence Against Women*. Ottawa: Ministry of Supply and Services Canada, March, 1996.
- Heffernan, K. "Eating Disorders and Weight Concern among Lesbians." *International Journal of Eating Disorders*, 19 (1996): 127-138.
- Henley, N.M., M. Miller & J.A. Beazley. "Syntax, Semantics, and Sexual Violence: Agency and the Passive Voice." *Journal of Language and Social Psychology*, 14, 1-2 (1995): 60-84.
- Hernandez, J. "The Concurrence of Eating Disorders with Histories of Child Abuse among Adolescents." *Journal of Child Sexual Abuse*, 4, 3 (1995): 73-85.

- Herscovici, C.R. & L. Bay "Favourable Outcome for Anorexia Nervosa Patients Treated in Argentina with a Family Approach." *Eating Disorders*, 4 (1996): 59-65.
- Herzog, D.B., M.B. Keller, P.W. Lavori & I.S. Bradburn. "Bulimia Nervosa in Adolescence." *Developmental and Behavioral Pediatrics*, 12, 3 (1991): 191-195.
- Herzog, D.B., J.E. Staley, S. Carmody, W.M. Robbins & B.A. van der Kolk. "Childhood Sexual Abuse in Anorexia Nervosa and Bulimia Nervosa: A Pilot Study." *Journal of the American Academy of Child and Adolescent Psychiatry*, 32 (1993): 962-966.
- Herzog, W., H.C. Deter, W. Fiehn & E. Petzold. "Medical Findings and Predictors of Long-Term Physical Outcome in Anorexia Nervosa: A Prospective 12-Year Follow-Up Study." *Psychological Medicine*, 27 (1997): 269-279.
- Hesse-Biber, S. "Eating Patterns and Disorders in a College Population: Are College Women's Eating Problems a New Phenomenon?" *Sex Roles*, 20 (1989): 71-89.
- Hill, J.P. & G.N. Holmbeck. "Familial Adaptation to Biological Change during Adolescence." In *Biological-Psychological Interactions in Early Adolescence*, ed. R.M. Lerner & T.T. Foch, 207-223. Hillsdale, NJ: Erlbaum, 1987.
- Hoek, H.W. "The Distribution of Eating Disorders." In *Eating Disorders and Obesity: A Comprehensive Handbook*, ed. K.D. Brownell & C.G. Fairburn, 207-211. New York: Guilford Press, 1995.
- Hof, S. van't & M. Nicolson. "The Rise and Fall of a Fact: The Increase in Anorexia Nervosa." *Sociology of Health and Illness*, 18 (1996): 581-608.
- Hsu, L.K. & S. Lee. "Is Weight Phobia Always Necessary for a Diagnosis of Anorexia Nervosa?" *American Journal of Psychiatry*, 150 (1993): 1466-1471.
- Hsu, L.K. & B. Zimmer. "Eating Disorders in Old Age." *International Journal of Eating Disorders*, 7 (1988): 133-138.
- Jiwani, Y. "Violence against Marginalized Girls." In *The Girl Child and Violence Prevention: A Review of Three Areas*, ed. Y. Jiwani & S. Moore with the assistance of S.M. Brown & B. Bunjun, 37-50. Vancouver: FREDA, 1998.
- Jiwani, Y., with the assistance of S. Moore & P. Kachuk. *Rural Women and Violence: A Study of Two Communities in British Columbia*. Report for the Department of Justice. Vancouver: FREDA, 1998.
- Joiner, G.W. & S. Kashubeck "Acculturation, Body Image, Self-Esteem, and Eating-Disorder Symptomatology in Adolescent Mexican American Women." *Psychology of Women Quarterly*, 20, 3 (1996): 419-435.
- Kaner, A., C.M. Bulik & P.F. Sullivan. "Abuse in Adult Relationships of Bulimic Women." *Journal of Interpersonal Violence*, 8 (1993): 52-63.
- Kennedy, M. "Patients Have Died Waiting, Doctors' Report Charges." *Ottawa Citizen*, August 20, 1998.

- Kinzl, J.F., C. Traweger, V. Guenther & W. Biebl "Family Background and Sexual Abuse Associated with Eating Disorders." *American Journal of Psychiatry*, 151, 8 (1994): 1127-1131.
- Koss, M.P. "Hidden Rape: Sexual Aggression and Victimization in a National Sample of Students in Higher Education." In *Violence in Dating Relationships: Emerging Social Issues*, ed. M.A. Pirog-Good & J.E. Stet, 143-168. New York: Praeger, 1989.
- Kraft, F. "Eating Disorders: Reducing the Risks." *Canadian Jewish News* (February 12, 1998): 57-58.
- Kuba, S.A. & S.G. Hanchey. "Reclaiming Women's Bodies: A Feminist Perspective on Eating Disorders." In *Feminist Perspectives on Addictions*, ed. Nan Van Den Bergh, 125-137. New York: Springer Publishing, 1991.
- Kufeldt, K., J. Baker, L. Bennett & R. Tite. *Looking After Children in Canada: Interim Draft Report*. St. John's, NF: Memorial University, 1998.
- LaBarbera, J.D. "Seductive Father-Daughter Relationships and Sex Roles in Women." *Sex Roles*, 11 (1984): 941-951.
- Lanzi, G., G. Rossi, U. Balottin, A. Citterio, G. Leonardi, T. Martelli, B. Seelsa, C. Tebaldi & C.A. Zambrino. "Disturbi del compartamento alimentare: Studio epidemiologico di 434 adolescenti." *Minerva Pediatrica*, 49, 6 (1997): 249-260.
- Larkin, J. & K. Popaleni. "Heterosexual Courtship Violence and Sexual Harassment: The Private and Public Control of Young Women." *Feminism and Psychology*, 4, 2 (1994): 213-227.
- LeBow, M.D. "Attitudes, Perceptions, and Practices of Canadian School Children towards Obesity." *Journal of Obesity and Weight Regulation*, 7, 1 (1988): 43-55.
- LeBow, M.D., D. Ness, P. Makarenko, T. Lam. "Attitudes, Perceptions and Practices of Canadian Teenagers about Obesity." *Journal of Obesity and Weight Regulation*, 8, 1 (1989): 53-65.
- Lawson, M.C. "Development of Body Build Stereotypes, Peer Ratings, and Self-Esteem in Australian Children." *Journal of Psychology*, 104, 1 (1980): 111-118.
- Lee, A.M. & S. Lee "Disordered Eating and Its Psychosocial Correlates among Chinese Adolescent Females in Hong Kong." *International Journal of Eating Disorders*, 20, 2 (1996): 177-183.
- Lee, S. "Self-Starvation in Context: Towards a Culturally Sensitive Understanding of Anorexia Nervosa." *Social Science and Medicine*, 41 (1995): 25-36.
- Lee, S., H.F.K. Chiu & C. Chen. "Anorexia Nervosa in Hong Kong: Why Not More Chinese?" *British Journal of Psychiatry*, 154 (1989): 683-688.
- le Grange, D., C.F. Telch & W.S. Agras. "Eating and General Psychopathology in a Sample of Caucasian and Ethnic Minority Subjects." *International Journal of Eating Disorders*, 21 (1997): 285-293.

- le Grange, D., C.F. Telch & J. Tibbs "Eating Attitudes and Behaviours in 1,435 South African Caucasian and Non-Caucasian College Students." *American Journal of Psychiatry*, 155 (1998): 250-254.
- Lieberman, M., L. Gauvin, W.M. Bukowski & D.R. White. "Interpersonal Influence and Disordered Eating Behaviors in Adolescent Girls: The Role of Peer Modeling, Social Reinforcement, and Body-Related Teasing." *Eating Behaviors*, 2 (2001): 215-236.
- Littlewood, R. "Psychopathology and Personal Agency: Modernity, Culture Change and Eating Disorders in South Asian Societies." *British Journal of Medical Psychology*, 68 (1995): 45-63.
- McCarthy, M. "Whose Body Is It Anyway? Pressures and Control for Women with Learning Disabilities." *Disability and Society*, 13 (1998): 557-574.
- McCreary Centre Society. *Adolescent Health Survey: Province of British Columbia*. Burnaby, BC: McCreary Centre Society, 1993.
- _____. *Adolescent Health Survey II: Province of British Columbia*. Burnaby, BC: McCreary Centre Society, 1998.
- _____. *Being Out: Lesbian, Gay, Bisexual & Transgender Youth in BC: An Adolescent Health Survey*. Burnaby, BC: McCreary Centre Society, 1999.
- McDonald, James A. "Building a Moral Community: Tsimshian Potlatching, Implicit Knowledge and Everyday Experiences." *Cultural Studies*, 9, 1 (1995): 125-144.
- MacDonald, John A. "Child Welfare and the Native Indian Peoples of Canada." *Windsor Yearbook of Access to Justice*, 5 (1985): 284-305.
- McKinley, N.M. & J.S. Hyde. "The Objectified Body Consciousness Scale: Development and Validation." *Psychology of Women Quarterly*, 20 (1996): 181-215.
- Malson, H. & J.M. Ussher. "Bloody Women: A Discourse Analysis of Amenorrhea as a Symptom of Anorexia Nervosa." *Feminism and Psychology*, 6 (1996): 505-521.
- Meister, J. "Women and Poverty." *Compass-Disabled Women and Poverty*. Winnipeg: Coalition of Provincial Organizations of the Handicapped, 1990.
- Mercer, S. Litch. *Not a Pretty Picture: An Exploratory Study of Violence against Women in High School Dating Relationships*. Toronto: Education Wife Assault, 1987.
- Miller, D.A.F., K. McCluskey-Fawcett & L.M. Irving. "The Relationship between Childhood Sexual Abuse and Subsequent Onset of Bulimia Nervosa." *Child Neglect and Abuse*, 17 (1993a): 305-314.
- _____. "Correlates of Bulimia Nervosa: Early Family Mealtime Experiences." *Adolescence*, 28 (1993b): 622-635.
- Miller, K.J. "Prevalence and Process of Disclosure of Childhood Sexual Abuse among Eating-Disordered Women." In *Sexual Abuse and Eating Disorders*, ed. Mark R. Schwartz & Leigh Cohn, 36-51. New York: Brunner/Mazel, 1996.

- Ministry of Health and Ministry Responsible for Seniors. *Policy Frameworks on Designated Populations*. Victoria, BC: Ministry of Health and Ministry Responsible for Seniors, April 1995.
- Mother's Report Call to Action Violence Against Midlife and Older Women, May 1994. Cited in *Elder Abuse*. Available online: <http://www.langara.bc.ca/vnc/elder.htm>
- Moyer, D.M., L. DiPietro, R.I. Berkowitz & A.J. Stunkard. "Childhood Sexual Abuse and Precursors of Binge Eating in an Adolescent Female Population." *International Journal of Eating Disorders*, 21 (1997): 23-30
- Mullen, P.E., J.L. Martin, J.C. Anderson, S.E. Romans & G.P. Herbison. "Childhood Sexual Abuse and Mental Health in Adult Life." *British Journal of Psychiatry*, 163 (1993): 721-732.
- Nagata, T., N. Kiriike, T. Iketani, Y. Kawarada & H. Tanaka. "History of Childhood Sexual or Physical Abuse in Japanese Patients with Eating Disorders: Relationship with Dissociation and Impulsive Behaviours." *Psychological Medicine*, 29 (1999): 935-942.
- National Aging Resource Center on Elder Abuse, 1990. Cited in *Elder Abuse*. Available online: <http://www.langara.bc.ca/vnc/elder.htm>
- National Clearinghouse on Family Violence. *Elder Abuse*. Ottawa: Health Canada, 1993.
- _____. *Youth and Violence*. Ottawa: Health Canada, 1997a. Available online: <http://www.hc-sc.gc.ca/hppb/familyviolence/pdfs/youthenglish.pdf>
- _____. *Child Sexual Abuse*. Ottawa: Health Canada, 1997b. Available online: <http://www.hc-sc.gc.ca/hppb/familyviolence/html/csaeng.html>
- _____. *Abuse of Older Adults in Institutions*. Ottawa: Health Canada, 2000a. Available online: http://www.hc-sc.gc.ca/hppb/familyviolence/html/older_adults%20_in_institutions/english/index.htm
- _____. *A Handbook Dealing with Woman Abuse and the Canadian Criminal Justice System: Guidelines for Physicians*. Ottawa: Health Canada, 2000b. Available online: <http://www.hc-sc.gc.ca/hppb/familyviolence/html/physician/english/index.htm>
- National Crime Prevention Council Canada. *Offender Profiles*, September 1995. Available online: <http://www.hc-sc.gc.ca/hppb/familyviolence/html/1offender.htm>
- Neumark-Sztainer, D., M. Story, N.H. Falkner, T. Beuhring & M.D. Resnick. "Disordered Eating among Adolescents with Chronic Illness and Disability: The Role of Family and other Social Factors." *Archives of Pediatric and Adolescent Medicine*, 152 (1998): 871-878.
- Neumark-Sztainer, D., M. Story, P.J. Hannan, T. Beuhring & M.D. Resnick. "Disordered Eating among Adolescents: Associations with Sexual/Physical Abuse and Other Familial/Psychosocial Factors." *International Journal of Eating Disorders*, 28 (2000): 249-258.

- Nicholson, S.D. & E. Ballance "Anorexia Nervosa in Later Life: An Overview." *Hospital Medicine*, 59 (1998): 268-272.
- Nielson, G.B., B. Lausch & P.H. Thomsen. "Three Cases of Severe Early-Onset Eating Disorder: Are They Cases of Anorexia Nervosa." *Psychopathology*, 30 (1997): 49-52.
- Offord, D.R., M.H. Boyle, D. Campbell, P. Goering, E. Lin, M. Wong & Y. Racine. "One-Year Prevalence of Psychiatric Disorder in Ontarians 15 to 64 Years of Age." *Canadian Journal of Psychiatry*, 41 (1996): 559-563.
- Oliver, K.K. & M.H. Thelen. "Children's Perceptions of Peer Influence on Eating Concerns." *Behavior Therapy*, 27 (1996): 25-39.
- Palmer, R.L. "Weight Concern Should Not Be a Necessary Criterion for the Eating Disorders: A Polemic." *International Journal of Eating Disorders*, 14 (1993): 459-465.
- Parry-Jones, B. & W.L. Parry-Jones. "History of Bulimia and Bulimia Nervosa." In *Eating Disorders and Obesity: A Comprehensive Handbook*, ed. K.D. Brownell & C.G. Fairburn, 145-150. New York: Guilford Press, 1995.
- Patterson, C. "Secondary Prevention of Elder Abuse." In *Canadian Task Force on the Periodic Health Examination*, by S.H. Woolf, R.N. Battista, G.M. Anderson, A.G. Logan & E.E. Wang, 922-929. Ottawa: Health Canada 1994.
- Paxton, S.J., H.K. Schutz, E.H. Wertheim & S.L. Muir. "Friendship Clique and Peer Influences on Body Image Concerns, Dietary Restraint, Extreme Weight-Loss Behaviors, and Binge Eating in Adolescent Girls." *Journal of Abnormal Psychology*, 108 (1999): 255-266.
- Pedersen, R. "Crazy Health Cuts Wasting Money – Top MD." *Edmonton Journal*, March 20, 1998.
- Peters, J.F. "Gender Socialization of Adolescents in the Home: Research and Discussion." *Adolescence*, 29 (1994): 913-934.
- Pike, K.M. & B.T. Walsh. "Ethnicity and Eating Disorders: Implications for Incidence and Treatment." *Psychopharmacology Bulletin*, 32 (1996): 265-274.
- Pope, H.G. & J.T. Hudson. "Is Childhood Sexual Abuse a Risk Factor for Bulimia Nervosa?" *American Journal of Psychiatry*, 149 (1992): 455-463.
- Pope, K.S. & V.A. Vetter. "Ethical Dilemmas Encountered by Members of the American Psychological Association: A National Survey." *American Psychologist*, 47 (1992): 397-411.
- Prince, R. "The Concept of Culture-Bound Syndromes: Anorexia Nervosa and Brain-Fag." *Social Science and Medicine*, 21, 2 (1985): 197-203.
- Pryor, D.W. "Fear of Stranger vs. Acquaintance Rape among College Women." Paper presented at the *American Sociological Association*, 1999.

- Razack, S. "From Consent to Responsibility, from Pity to Respect: Subtexts in Cases of Sexual Violence Involving Girls and Women with Developmental Disabilities." *Law and Social Inquiry*, 19 (1994): 891-922.
- Reid, P. & G. Finchilescu. "The Disempowering Effects of Media Violence against Women on College Women." *Psychology of Women Quarterly*, 19, 3 (1995): 397-411.
- Reitsma-Street, M. "Justice for Canadian Girls: A 1990s Update." *Canadian Journal of Criminology*, 41, 3 (July 1999): 335-364.
- Reto, C.S. "Childhood Maltreatment, Dissociation, and Bulimic Symptomatology." *Dissertation Abstracts International*, 58, 12-B (1998): 6823.
- Reto, C.S., C.J. Dalenberg & M.T. Coe. "Dissociation and Physical Abuse as Predictors of Bulimic Symptomatology and Impulse Dysregulation." In *Sexual Abuse and Eating Disorders*, ed. Mark R. Schwartz & Leigh Cohn, 52-67. New York: Brunner/Mazel, 1996.
- Ristock, J. "Beyond Ideologies: Understanding Violence in Lesbian Relationships." *Canadian Woman Studies*, 12 (1991): 74-79.
- Ritter, M.B. "Mothers, Daughters, Culture and Criticism: An Examination of Eating Problems in Adolescent Girls." *Dissertation Abstracts International*, 58, 9-B (1998): 5138.
- Root, M.P.P. "Disordered Eating in Women of Colour." *Sex Roles*, 22 (1990): 525-536.
- Root, M.P.P. & P. Fallon "The Incidence of Victimization Experiences in a Bulimic Sample." *Journal of Interpersonal Violence*, 3 (1988): 161-173.
- Rorty, M. & J. Yager "Speculations on the Role of Childhood Abuse in the Development of Eating Disorders among Women." In *Sexual Abuse and Eating Disorders*, ed. Mark R. Schwartz & Leigh Cohn, 23-35. New York: Brunner/Mazel, 1996.
- Rosen, L.W., C.L. Shafer, G.M. Dummer, L.K. Cross, G.W. Deuman & S.R. Malmberg. "Prevalence of Pathogenic Weight-Control Behaviours among Native American Women and Girls." *International Journal of Eating Disorders*, 7 (1988): 807-811.
- Roush, D.J. "A Qualitative Study of Sex Abuse Survivors' Experience of Confronting the Perpetrator." *Dissertation Abstracts International*, 59, 11-A (1999): 4294.
- Ryn, Z. "Between Life and Death: Experiences of Concentration Camp Musselmen During the Holocaust." *Genetic, Social, and General Psychology Monographs*, 116 (1990): 7-19.
- Samis, Stephen M. "An Injury to One is an Injury to All': Heterosexism, Homophobia, and Anti-Gay/Lesbian Violence in Greater Vancouver." M.A. Thesis. Simon Fraser University, 1995.
- Schaaf, K.K. & T.R. McCanne. "Childhood Abuse, Body Image Disturbance, and Eating Disorders." *Child Abuse and Neglect*, 18 (1994): 607-615.

- Schmidt, U., H. Humfress & J. Treasure. "The Role of General Family Environment and Sexual and Physical Abuse in the Origins of Eating Disorders." *European Eating Disorders Review*, 5 (1997): 184-207.
- Schmidt, U., J. Tiller, M. Blanchard, B. Andrews & J. Treasure "Is There a Specific Trauma Precipitating Anorexia Nervosa?" *Psychological Medicine*, 27 (1997): 523-530.
- Schmidt, U., J. Tiller & J. Treasure "Setting the Scene for Eating Disorders: Childhood Care, Classification and Course of Illness." *Psychological Medicine*, 23 (1993): 663-672.
- Senior Women Against Abuse Collective. *Elder Women Speak Out on Abuse*. Winnipeg: Manitoba Council on Aging, 1989.
- Siever, M.D. "The Perils of Sexual Objectification: Sexual Orientation, Gender, and Socioculturally Acquired Vulnerability to Body Dissatisfaction and Eating Disorders." In *Gay and Lesbian Mental Health: A Sourcebook for Practitioners*, ed. Christopher J. Alexander, 223-247. New York: Harrington Park Press, 1996.
- Simkin, R.J. "Lesbians Face Unique Health Care Problems." *Canadian Medical Association Journal*, 145 (1991): 1620-1623.
- _____. "Not All Your Patients are Straight." *Canadian Medical Association Journal*, 159 (1998): 370-375.
- Smith, J.E. & J. Krejci "Minorities Join the Majority: Eating Disturbances among Hispanic and Native American Youth." *International Journal of Eating Disorders*, 10, 2 (1991): 179-186.
- Smith, J.E., V.A. Waldorf & D.L. Trembath. "Single White Male Looking for Thin, Very Attractive ..." *Sex Roles*, 23 (1990): 675-685.
- Smith, S. "Women's Experiences of Victimizing Sexualization, Part 1: Responses Related to Abuse and Home and Family Environment." *Issues in Mental Health Nursing*, 18 (1997): 395-416.
- Snow, J.T. & M.B. Harris. "Disordered Eating in South-Western Pueblo Indians and Hispanics." *Journal of Adolescence*, 12 (1989): 329-336.
- Statistics Canada. *Violence Against Women Survey*. Victoria, BC: Ministry of Women's Equality, 1993.
- _____. *Women in Canada: A Statistical Report*. Ottawa: Statistics Canada, 1995.
- Striegel-Moore, R.H. "A Feminist Perspective on the Etiology of Eating Disorders." In *Eating Disorders and Obesity: A Comprehensive Handbook*, ed. K.D. Brownell & C.G. Fairburn: 224-229. New York: Guilford Press, 1995.
- Stump, C.W. "Women's Eating Attitudes and Body Image in Coed and Single Sex High Schools: An Exploration of Social Contagion." *Dissertation Abstracts International*, 55, 10-B (1995): 4615.

- Thompson, B.W. "A Way Outa No Way: Eating Problems among African-American, Latina, and White Women." *Gender and Society*, 6, 4 (1992): 564-561.
- _____. *A Hunger So Wide and So Deep: A Multiracial View of Women's Eating Problems*. Minneapolis, MN: University of Minnesota Press, 1994.
- Tsiantas, G. & R.M. King. "Similarities in Body Image in Sisters: The Role of Sociocultural Internalization and Social Comparison." *Eating Disorders: The Journal of Treatment and Prevention*, 9 (2001): 141-158.
- Twamley, E.W. & M.C. Davis. "The Sociocultural Model of Eating Disturbance in Young Women: The Effects of Personal Attributes and Family Environment." *Journal of Social and Clinical Psychology*, 18 (1999): 467-489.
- Vincent, M.A. & M.P. McCabe. "Gender Differences among Adolescents in Family, and Peer Influences on Body Dissatisfaction, Weight Loss, and Binge Eating Behaviors." *Journal of Youth and Adolescence*, 29 (2000): 205-221.
- Waller, G., C. Halek & A.H. Crisp. "Sexual Abuse as a Factor in Anorexia Nervosa: Evidence from Two Separate Case Series." *Journal of Psychosomatic Research*, 37 (1993): 873-879.
- Waller, G., A. Ruddock & S. Cureton. "Cognitive Correlates of Reported Sexual Abuse in Eating." *Journal of Interpersonal Violence*, 10 (1995): 176-187.
- Watson, L.A. "Mirror, Mirror on the Wall ... : An Exploratory Study of Physical Disability and Women's Expressed Body Image." *Dissertation Abstracts International*, 59, 9-B (1999): 5173.
- Websdale, N. "Rural Woman Abuse: The Voices of Kentucky Women." *Violence Against Women*, 1 (1995a): 309-338.
- _____. "An Ethnographic Assessment of the Policing of Domestic Violence in Rural Kentucky." *Social Justice*, 22 (1995b): 102-122.
- _____. *Rural Woman Battering and the Justice System: An Ethnography*. Thousand Oaks, CA: Sage Series on Violence Against Women, 1998.
- Weiner, E.J. & L. Stephens. "Sexual Barrier Weight: A New Approach." In *Sexual Abuse and Eating Disorders*, ed. Mark R. Schwartz & Leigh Cohn, 68-87. New York: Brunner/Mazel, 1996.
- Welch, S.L., H.A. Doll & C.G. Fairburn. "Life Events and the Onset of Bulimia Nervosa: A Controlled Study." *Psychological Medicine*, 27 (1997): 515-522.
- Welch, S.L. & C.G. Fairburn. "Childhood Sexual and Physical Abuse as Risk Factors for the Development of Bulimia Nervosa: A Community-Based Case Control Study." *Child Abuse and Neglect*, 20 (1996): 633-642.
- Wiederman, M.W. "Eating Disorders in an 86-Year-Old Female." *Clinical Gerontologist*, 16, 2 (1996): 77-79.
- Williams, H.J., H.L. Wagner & R.M. Calam. "Eating Attitudes in Survivors of Unwanted Sexual Experiences." *British Journal of Clinical Psychology*, 31 (1992): 203-206.

- Williamson, L. "Eating Disorders and the Cultural Forces behind the Drive for Thinness: Are African American Women Really Protected?" *Social Work in Health Care*, 28 (1998): 61-73.
- Wonderlich, S.A., T.Y. Brewerton, Z. Jolic, B.S. Dansky & D.W. Abbott. "Relationship of Childhood Sexual Abuse and Eating Disorders." *Journal of the American Academy of Child and Adolescent Psychiatry*, 36, 8 (1997): 1107-1115.
- Woodside, D.B. "A Review of Anorexia Nervosa and Bulimia Nervosa." *Current Problems in Pediatrics*, 25 (1995): 67-89.
- Yoder, J.D. *Women and Gender: Transforming Psychology*. Upper Saddle River, NJ: Prentice Hall, 1999.
- Zani, B. "Male and Female Patterns in the Discovery of Sexuality During Adolescence." *Journal of Adolescence*, 14 (1991): 163-178.
- Zerbe, K.J. "Eating Disorders in the 1990s: Clinical Challenges and Treatment Implications." *Bulletin of the Menninger Clinic*, 56 (1992): 167-187.

APPENDIX A: THE ADVISORY PANEL

Sheila Baxter

Author and Anti-Poverty Activist
Vancouver, BC

Merryl Bear

National Eating Disorders Information Centre (NEDIC)
Toronto, ON

Marion Fallding

Counsellor
Chilliwack Mental Health
Chilliwack, BC

Sandy Friedman

Author and Researcher
Vancouver, BC

Dr. Sharon Fuller

Department of Anthropology and Sociology
Chair, Cultural and Media Studies Interdisciplinary Group
Green College, UBC
Vancouver, BC

Dr. John Gainer

Psychologist
Cape Breton Healthcare Complex
Sydney, NS

Abigail Idlout

Sexual Abuse Specialist
Baffin Region Health and Social Services Board
Iqaluit, NT

Laura Quilici

Co-coordinator, Vancouver Coordination Program on Violence against Women in
Relationships, Battered Women's Support Services
Vancouver, BC

Françoise Mathieu

Counsellor
Queen's University Student Counselling Service
Kingston, ON

Raine McKay

Director

Association for Awareness and Networking Around Disordered Eating (ANAD)
Vancouver, BC

Janice Williams

Director

The Centre for Recovering Anorexic:bulimic Disordered Lives Entrust (CRADLE)
Lethbridge, AB

Shelley Wilson

Pilot Project Coordinator, Girls in the 90's
Public Health Services
Yarmouth, NS

FREDA PARTICIPANTS

Kelly D'Aoust

Researcher, FREDA

Dr. Yasmin Jiwani

Executive Director, FREDA

Dr. Shelley Moore

Research Associate, FREDA
Author of "Swallowing the Hurt"

Donna Robertson

Research Coordinator, FREDA

APPENDIX B: A LIST OF PROGRAMMES

This appendix lists programmes providing services for disordered eating. No individual programme listed is necessarily recommended or endorsed by FREDA or by Health Canada. An extensive directory of transition houses, emergency shelters, second stage housing and safe networks across Canada has been published in *Transition Houses and Shelters for Abused Women in Canada* (2000), available through the National Clearinghouse on Family Violence, Health Canada.

BRITISH COLUMBIA

Chilliwack Mental Health
274 - 45470 Menholm Road
Chilliwack, BC, V2P 1M2
Tel: 604-795-8375

Eating Disorder Outreach Program
St. Joseph's Hospital
2137 Comox Avenue
Comox, BC, V9M 1P2
Tel: 250-339-1576
Fax: 250-339-1439

East Kootenay Eating Disorder Clinic
Cranbrook Regional Hospital
13 - 24 Avenue, North
Cranbrook, BC, V1C 3H9
Tel: 250-489-6416
Fax: 250-426-5285

Tri-Cities Mental Health BC
2232 Elgin Avenue
Port Coquitlam, BC, V3C 3B2
Tel: 604-941-3471
Fax: 604-660-9805

Ministry of Children and Family
Development, Tri-Cities BC
300 - 3003 St. John's Street
Port Moody, BC, V3H 2C4
Tel: 604-469-7600
Fax: 604-469-7601

HUGS, Cranbrook Regional Hospital
13 - 24th Avenue North
Cranbrook, BC, V1C 3H9
Tel: 250-426-5281
Fax: 250-426-6262

Child & Youth Team
Canadian Mental Health Association
205 - 149 Ingram Street
Duncan, BC, V9L 1N8
Tel: 250-746-5521
Fax: 250-748-2606

Mental Health
PO Box 2126
Invermere, BC, V0A 1K0
Tel: 250-342-4295
Fax: 250-342-4322

Kamloops Community Eating Disorder
Program
Thompson Health Region
519 Columbia Street
Kamloops, BC, V2C 2T8
Tel: 250-828-4438
or 250-828-4143
Fax: 250-828-4990

Langley Memorial Hospital
22051 Fraser Hwy
Langley, BC, V3A 4H4
Tel: 604-533-6493

Nanaimo Family Life Association
1070 Townsite Road
Nanaimo, BC, V9S 1M6
Tel: 250-754-3331
Fax: 250-753-0268

Associated Family and Community
Support Services, Ltd.
2033 Sanders Road
Nanose Bay, BC, V9P 9C2
Tel: 250-248-0076
Fax: 250-468-9182

Eating Disorders Self Help Group
Kootenay Lake Regional Hospital
333 Victoria Street
Nelson, BC, V1L 4K3
Tel: 250-354-6321
Fax: 250-354-6320

North Shore Mental Health BC
209 - 267 West Esplanade
North Vancouver, BC, V7M 1A5
Tel: 604-660-1273
Fax: 604-660-3108

ANAD North Vancouver Support Group
North Vancouver City Library
121 West 14 Street
North Vancouver, BC, V7M 1N9
Tel: 604-998-3450
Fax: 604-983-3624

Tri-Cities Mental Health
Eating Disorders Program
2232 Elgin Avenue
Port Coquitlam, BC, V3C 3B2
Tel: 604-941-3471
Fax: 604-660-9805

Child & Youth Mental Health Services,
300 - 3003 St. John's Street
Port Moody, BC, V3H 2C4
Tel: 604-469-7600
Fax: 604-469-7601

HUGS, c/o Heart & Stroke Foundation
106 - 490 Quebec Street
Prince George, BC, V2L 5N5
Tel: 250-562-8611
Fax: 250-562-8614

Prince George Eating Disorder Clinic
Northern Interior Health Unit
1444 Edmonton Street
Prince George, BC, V2M 6W5
Tel: 250-565-7479
Fax: 250-565-7416

Shuswap Lake General Hospital
PO Box 520 STN MAIN
Salmon Arm, BC, V1E 4N6
Tel: 250-833-3636 Ext. 259
Fax: 250-833-3602

Sunshine Coast Mental Health Services,
St. Mary's Hospital
PO Box 949
5544 Highway 101
Sechelt, BC, V0N 3A0
Tel: 604-885-6101
Fax: 604-885-5842

Bulkley Valley District Hospital
PO Box 370
Smithers, BC, V0J 2N0
Tel: 604-847-2611 Ext. 255

National Institute for Compulsive Eaters
2467 - 127B Street
Surrey, BC, V4A 8N8
Tel: 604-873-6423
Fax: 604-536-6423

Surrey Central Mental Health
110 - 7525 King George Highway
Surrey, BC, V3W 5A8
Tel: 604-543-5660
Fax: 604-543-5699

FAIR Family & Individual Resources
860 Eldorado Street, PO Box 153
Trail, BC, V1R 4L5
Tel: 250-364-2326 or 250-368-3311

Association for Awareness and
Networking around Disordered Eating
109 - 2040 West 12 Avenue
Vancouver, BC, V6J 2G2
Tel: 604-739-2070
Fax: 604-730-2843

ANAD East Vancouver Support Group,
Mount Pleasant Neighbourhood House
4 - 800 East Broadway
Vancouver, BC, V5T 1Y1
Tel: 604-879-8208

ANAD Kitsilano Support Group
West Side Family Place
2819 West 11th Avenue
Vancouver, BC, V6K 2M2
Tel: 604-738-2819
Fax: 604-738-2850

Healthy Attitudes Program
Vancouver/Richmond Health Board
6405 Knight Street
Vancouver, BC, V5P 2V9
Tel: 604-321-6151
Fax: 604-321-2947

Supporting the Struggle Against
Anorexia Nervosa (SUSTAAN)
PO Box 29105 RPO Delamont
Vancouver, BC, V6J 5C2
Tel: 604-734-0006

BC Coalition to End Disordered Eating
Eating Disorders Program for Children &
Adolescents, BC Children's Hospital
4480 Oak Street
Vancouver, BC, V6H 3V4
Tel: 604-875-2200
Fax: 604-875-2271

Nutrition Clinic at the Children's Centre,
Mount St. Joseph's Hospital
3080 Prince Edward Street
Vancouver, BC, V5T 3N4
Tel: 604-877-8551

Eating Disorder Service
University Hospital – UBC Site
2211 Wesbrook Mall
Vancouver, BC, V6T 2B5
Tel: 604-822-2415

Eating Disorder Resource Centre of BC
St. Paul's Hospital
1081 Burrard Street
Vancouver, BC, V6Z 1Y6
Tel: 604-631-5313 or 1-800-665-1822
Fax: 604-631-5461

Vancouver Anti-anorexia/ Anti-bulimia
League (VAAABL)
306 - 1212 West Broadway
Vancouver, BC, V6H 3V1
Tel: 604-731-7304
Fax: 604-730-1015

Deliciosa! Nutrition Counselling
3675 West 16th Avenue
Vancouver, BC, V6R 3C3
Tel: 604-225-0505
Fax: 604-225-0555

What Are You Hungry For?
3675 West 16th Avenue
Vancouver, BC, V6R 3C3
Tel: 604-225-0505
Fax: 604-225-0555

Vernon Eating Disorder Program for
Youth and Adults/The People Place
303 - 3402 27th Avenue
Vernon, BC, V1T 1S1
Tel: 250-542-7111
Fax: 250-542-7111

Vernon & Area
Eating Disorders Association
Vernon, BC
Tel: 250-542-1388

BC Eating Disorders Association
526 Michigan Street
Victoria, BC, V8V 1S2
Tel: 250-383-2755
Fax: 250-383-5518

Eros House for Creative Recovery.
628 Bryden Court
Victoria, BC, V9A 4Y5
Tel: 250-361-4848

Ministry of Children and Family
Development
302 - 2955 Jutland Road
Victoria, BC, V8T 5J9
Tel: 250-387-0000
Fax: 250-387-0002

Ministry of Children and Family
Development, North Shore
230 - 1425 Marine Drive
West Vancouver, BC, V7T 1B9
Tel: 604-981-0165
Fax: 604-926-5835

Outpatient Eating Disorders Program
Williams Lake Mental Health Centre
487 Borland Street
Williams Lake, BC, V2G 1R9
Tel: 250-398-4465

YUKON

Yukon Mental Health Services
PO Box 2703
4 Hospital Road
Whitehorse, YT, Y1A 3H8
Tel: 867-667 8346
Fax: 867-667-8372

NORTHWEST TERRITORIES

HUGS, H.H. Williams Memorial Hospital
3 Gaetz Drive
Hay River, NT, X0E 0R8
Tel: 867-874-6512 Ext. 140
Fax: 867-874-3377

Baffin Region Health and Social Services Board
Community Services
Bag 200
Iqaluit, NT, X0A 0H0
Tel: 867-979-7680

ALBERTA

Eating Disorders Co-ordinator
Calgary Regional Health Authority
1509 Centre Street S., 4th Floor
Calgary, AB, T2G 2G6
Tel: 403-303-6002
Fax: 403-232-6153

Eating Disorders Program
Outpatient Services
Calgary Counselling Centre
200 - 940 6 Avenue SW
Calgary, AB, T2P 3T1
Tel: 403-265 4980
Fax: 403-256-8886

HUGS, Coronation Health Centre
PO Box 250, Mailbag 500
Coronation, AB, T0C 1C0
Tel: 403-578-3803
Fax: 403-578-3474

Still Waters Counselling Service
PO Box 1126
Crossfield, AB, T0M 0S0
Tel: 403-703-3743
Fax: 403-337-2136

HUGS
Drumheller District Health Services
625 Riverside Drive East
Drumheller, AB, T0J 0Y0
Tel: 403-820-7213
Fax: 403-823-5076

University of Alberta Hospital Site
Subunit 4F4
8440 - 112 Street
Edmonton, AB, T6G 2B7
Tel: 780-492-6114
Fax: 780-492-1310

Elizabeth Seton Community Partners
c/o Community Services – Clairview Site
600A Hermitage Road
Edmonton, AB, T5A 4N2
Tel: 780-496-5868
Fax: 780-496-5881

HUGS, c/o Kinsmen Sports Centre
9100 Walterdale Hill NW
Edmonton, AB, T6E 2V3
Tel: 780-496-7300

HUGS
Stetsko Mayne Nutrition Consulting
7802 Mission Heights Drive
Grande Prairie, AB, T8W 1Y2
Tel: 780-538-1275
Fax: 780-532-7034

HUGS, Harry Collinge High School
158 Sunwapta Drive
Hinton, AB, T7V 1T7
Tel: 780-865-3714
Fax: 780-865-5011

The Centre for Recovering
Anorexic:bulimic Disordered Lives
Entrust (CRADLE)
50 Ryerson Bay West
Lethbridge, AB, T1K 4P4
Tel: 403-381-8544

SASKATCHEWAN

HUGS

Pipestone Health District Mental Health
PO Box 970
Grenfell, SK, S0G 2B0
Tel: 306-697-3577
Fax: 306-697-2686

Girls in the 90's, Pasquia Health Unit
PO Box 1075
Hudson Bay, SK, S0E 0Y0
Tel: 306-865-3277
Fax: 306-865-2660

North Central Health District Mental
Health Services
Melfort Hospital
PO Box 1480
Melfort, SK, S0E 1A0
Tel: 306-752-8767
Fax: 306-752-8711

BridgePoint Center for Eating Disorders
PO Box 190
Milden, SK, S0L 2L0
Tel: 306-935-2240
Fax: 306-935-2241

Moose Jaw Mental Health Clinic
455 Fairford Street East
Moose Jaw, SK, S6H 1H3
Tel: 306-691-6464
Fax: 306-691-6461

HUGS

1071 River Street East
Prince Albert, SK, S6V 7N6
Tel: 306-764-5820

Saskatchewan Provincial Consultant for
Eating Disorders
Saskatchewan Ministry of Health
3475 Albert Street
Regina, SK, S4S 6X6
Tel: 306-655-6673

Inter-Agency Committee for the
Prevention and Management of Eating
Disorders
350 Cheadle Street West
Swift Current, SK, S9H 4G3
Tel: 306-778-5250
Fax: 306-778-5408
Contact: Krista Olson, Social Worker;
Cathy Knox, Public Health Nutritionist

MANITOBA

Westwind Eating Disorder Recovery
458 - 14th Street
Brandon, MB, R7A 4T3
Tel: 204-728-2499

HUGS
Box 102A, RR 3
Portage La Prairie, MB, R1N 3A3
Tel: 204-428-3432
Fax: 204-428-5072

Women's Health Clinic
419 Graham Avenue, 3rd Floor
Winnipeg, MB, R3C 0M3
Tel: 204-947-1517
Fax: 204-943-3844

HUGS
518 - 1281 Grant Avenue
Winnipeg, MB, R3M 1Z6
Tel: 204-478 4847
Fax: 204-488 2169

Winnipeg Eating Disorder Clinic
Health Sciences Centre
771 Bannatyne Avenue
Winnipeg, MB, R3E 3N4
Tel: 204-787-3345 or 204-787-3482

ONTARIO

Body Image Coalition of Peel
180 B Sandalwood Parkway, E.
Brampton, ON, L6Z 4N1
Tel: 905-791-7800 Ext. 7694

HUGS
22 Westgate Walk
Brampton, ON, L6Y 3H4
Tel: 905-453-5590

New Attitudes and New Directions
135 McHardy Court
Brampton, ON, L6Y 1H7
Tel: 905-796-3474

Pediatric Program
Peel Memorial Hospital
20 Lynch Street
Brampton, ON, L6W 2Z8
Tel: 905-796-4066 Ext. 4010

Brant Community Mental Health Centre
408 - 760 Brant Street, Level 2
Burlington, ON, L7R 4B7
Tel: 905-631-1939
Fax: 905-631-0513

Eating Disorders Program
Joseph Brant Memorial Hospital
1230 N. Shore Blvd., E.
Burlington, ON, L7R 4C4
Tel: 905-632-3730

The Wellness Centre, Inc.
PO Box 364
Campbellville, ON, L0P 1B0
Tel: 905-854-2390

HUGS
287 Campus Parkway
Chatham, ON, N7L 4V7
Tel: 519-351-4292

HUGS Nutrition Counselling
4 Erinwood Drive
Erin, ON, N0B 1T0
Tel: 519-833-0843
Fax: 519-824-9233

Inpatient Eating Disorders Program
Homewood Health Centre
150 Delhi Street
Guelph, ON, N1E 6K9
Tel: 519-824-1762

Childrens Exercise and Nutrition Centre
Chedoke McMaster Hospital
Sanatorium Road, PO Box 2000
Hamilton, ON, L8N 3Z5
Tel: 905-521-7967
Fax: 905-385-5033

North Kingston Community Health
400 Elliot Avenue
Kingston, ON, K7K 6M9
Tel: 613-542-2813
Fax: 613-542-5486

Kingston Psychiatric Hospital (KPH)
PO Box 603 STN MAIN
752 King Street West
Kingston, ON, K7L 4X3
Tel: 613-546-1101
Fax: 613-548-5588
<http://meds.queensu.ca/kph/>

Beechgrove Children's Centre
PO Box 7777
Kingston, ON, K7L 5H1
Tel: 613-549-5600

Youth Crisis Service
Kingston, ON
Tel: 613-548-1155

Hotel Dieu Hospital Eating Disorders
Program, Hotel Dieu Hospital
166 Brock Street
Kingston, ON, K7L 5G2
Tel: 613-544-3310; 613-548-6121

KGH Outpatient Eating Disorders Clinic
72 Barrie Street
Kingston, ON, K7I 3J7
Tel: 613-548-6121

Kingston Community Counselling Centre
417 Bagot Street
Kingston, ON, K7K 9Z9
Tel: 613-549-7850
Fax: 613-544-8138

Anorexia Nervosa and Bulimia
Association (ANAB)
767 Bayridge Drive
PO Box 20058
Kingston, ON, K7P 1C0
Tel: 613-547-3684
<http://www.phe.queensu.ca/anab/>

Student Health Services, Queen's
University, St Lawrence Building
Kingston, ON, K7L 3N6
Tel: 613-533-2893
Fax: 613-533-6740

HUGS
PO Box 670 STN MAIN
Lindsay, ON, K9V 4W9
Tel: 705-454 9818
Fax: 705-454 9837

HUGS
PO Box 1046 STN MAIN
Lively, ON, P3Y 1M8
Tel: 705-692 0720

Eating Disorders Association of London
Victoria Family Medical Centre
60 Chesley Avenue
London, ON, N5Z 2C1
Tel: 519-433-8424
Fax: 519-433-2244

HUGS, Long Life-Style Consulting
PO Box 532
Maxville, ON, K0C 1T0
Tel: 613-527 3377
Fax: 613-527 3377

HUGS
1703 Kelsey Court
Mississauga, ON, L5L 3J8
Tel: 905-291 7573
Fax: 905-607 5420

HUGS, Health Source Associates
2550 Argentia Road
Mississauga, ON, L5N 5R1
Tel: 905-814 0448
Fax: 905-814 0448

Mississauga Community Health Nursing
Peel Health
3038 Hurontario
Mississauga, ON, L5B 3B9
Tel: 905-791-7800 Ext. 7401

Trillium Health Centre
Mississauga Hospital
100 Queensway West
Mississauga, ON, L5B 1B8
Tel: 905-848-7100
Fax: 905-848-7592

HUGS
185 Napier Street
Mitchell, ON, N0K 1N0
Tel: 519-348 4293
Fax: 519-348 4293

Eating Disorders Recovery Group
4 Bruno Street
Naughton, ON, P0M 2M0
Tel: 705-692-0442

HUGS, Nutrition Consultants Ottawa
91 Beaver Ridge
Nepean, ON, K2E 6E5
Tel: 613-224 5685
Fax: 613-723 9173

Jack Knight, Private Practice, Box 156
1100 Gorham Street, Suite 11B
Newmarket, ON, L3Y 7V1
Tel: 905-953-5685; 905-476-2880
Fax: 905-476-2880

HUGS, Perfect Balance Canada
2006 - 7 Bishop Avenue
North York, ON, M2M 4J4
Tel: 416-250 6658
Fax: 416-733 4719

Child & Adolescent Psychiatry Program
Oakville Trafalgar Memorial Hospital,
327 Reynolds Street
Oakville, ON, L6L 3L7
Tel: 905-338-4134

Bureau des services à la jeunesse
d'Ottawa-Carleton
1338 ½ Wellington Street
Ottawa, ON, K1Y 3B7
Tel: 613-729-1000
Fax: 613-729-1918
<http://www.ysb.on.ca/>

Department of Psychiatry
Ottawa General Hospital
501 Smyth Road, Box 400
Ottawa, ON, K1H 8L6
Tel: 613-737-8010
Fax: 613-739-9980

HUGS, OASIS Institute
R.R.3 - 1803
Prescott, ON, K0E 1T0
Tel: 613-657-4688
Fax: 613-925-4537

Eating Disorders Recovery Group
206 - 111 Elm Street
Sudbury, ON, P3C 1T3
Tel: 705-692-0442
<http://www.mirror-mirror.org/eatdis.htm>

HUGS, Diet Enders
23 Oriah Court
Thornhill, ON, L4J 8B3
Tel: 905-764 5935

Hospital for Sick Children
555 University Avenue
Toronto, ON, M5G 1X8
Tel: 416-813-7195
Fax: 416-813-7867
<http://www.sickkids.on.ca>

Ontario Centre for Adolescent Eating
Disorders
210 - 2 Gloucester Street
Toronto, ON, M4Y 1L5
Tel: 416-944-2693
Fax: 416-813-5560

Bellwood Health Services, Inc.
1020 McNicoll Avenue
Toronto, ON, M1W 2J6
Tel: 416-495-0926 & 1-800-387-6198
Fax: 416-495-7943
<http://www.bellwood.ca>

Anorexia Bulimia Family Support Group
783 Windermere Avenue
Toronto, ON, M6S 3M5
Tel: 416-766-8134
Fax: 416-762-5642

Nutritional Eating Disorder Clinic
1206 - 4950 Yonge Street
Toronto, ON, M2W 6K1
Tel: 416-229-6656 or 416-498-4921

Brief Psychotherapy Centre for Women
1806 - 2 Carlton Street
Toronto, ON, M5B 1J3
Tel: 416-591-2000

National Eating Disorders Information
Centre (NEDIC)
CW 1-211, 200 Elizabeth Street
Toronto, ON, M5G 2C4
Tel: 416-340-4156
Fax: 416-340-4736
<http://www.nedic.ca/>

Toronto Hospital
Program for Eating Disorders
EN 8-231, 200 Elizabeth Street
Toronto, ON, M5G 2C4
Tel: 416-340-3041
Fax: 416-340-4198

Ontario Community Outreach Program
for Eating Disorders
Toronto General Hospital
CCRW 2-828, 101 College Street
Toronto, ON, M5G 1L7
Tel: 416-340-4051

New Realities Eating Disorder Recovery
103 - 62 Charles Street East
Toronto, ON, M4Y 1T1
Tel: 416-921-9670; or 905-763-0660
<http://www.newrealitiescan.com>

Sheena's Place
87 Spadina Road
Toronto, ON, M5R 2T1
Tel: 416-927-8900
Fax: 416-927-8844

HUGS, The Wellington Club
111 Wellington Street West
Toronto, ON, M5J 2S6
Tel: 416-362 2582
Fax: 416-362 1373

Sandra Edwards, Private Practice
308 - 10 Unionville Gate
Unionville, ON, L3R 0W7
Tel: 905-479-0869 (office)
Fax: 905-946-1431

Body Balance Total Nutrition Care
105 University Avenue East
Waterloo, ON, N2J 2W1
Tel: 519-747-1848
Fax: 519-747-1848

HUGS
371 Prince of Wales Drive
Whitby, ON, L1N 6M8
Tel: 905-668 6831 Ext. 1336
Fax: 905-668 8279

Bulimia Anorexia Nervosa Association
(BANA)
300 Cabana Road East
Windsor, ON, N9G 1A3
Tel: 519-969-2112; Fax: 519-969-0227
<http://www.bana.ca>

Vitanova Foundation
6299 Rutherford Road
Woodbridge, ON, L4L 1A7
Tel: 905-850-3690

QUÉBEC

Centre de traitement des désordres
alimentaires du Québec
8149, rue du Mistral, Bur. 201
Charny, QC, G6X 1G5
Tel: 418-832-0574

Groupe de soutien et d'entraide Anorexie
et Boulimie de Granby
315, rue Cartier
Granby, QC, J2G 5A9
Tel: 450-372-2098
Fax: 450-372-0406

Le centre de thérapie du comportement
L'Hôpital Général de Montréal
1650 Cedar Avenue
Montréal, QC, H3G 1A4
Tel: 514-934-8034

Le collectif action alternative en obésité
7378 rue Lajeunesse, bur. 210
Montréal, QC, H2R 2H8
Tel: 514-270-3779
<http://www.multimania.com/caao/>

Outremangeurs Anonymes
434, rue De L'Église
Montréal, QC, H4G 2M4
Tel: 514-490-1939

Service de médecine à l'adolescence,
L'hôpital Sainte-Justine
3175, chemin de la Côte Ste-Catherine
Montréal, QC, H3T 1C5
Tel: 514-345-4721

Service de santé aux étudiants,
l'Université McGill
3600 McTavish
Montréal, QC, H3A 2T5
Tel: 514-392-5119

Anorexia & Bulimia Foundation of
Québec (ANAB)
114 Donegani Blvd.
Pointe Claire, QC, H9R 2W3
Tel: 514-630-0907
Fax: 514-630-1225
<http://www.generation.net/~anebque>

Clinique St-Amour
1100, Boul de la Rive-sud, Suite 120
St. Romuald, QC, G6W 5M6
Tel: 418-834-9825

Douglas Hospital, Eating Disorder Unit
6605 LaSalle Blvd.
Verdun, QC, H4H 1R3
Tel: 514-761-6131 Ext. 22894
Fax: 514-761-8885

NEW BRUNSWICK

HUGS

PO Box 3365 STN B
Fredericton, NB, E3A 5H2
Tel: 506-458 9285

HUGS

158 Burpee Street
Fredericton, NB, E3A 1M5
Tel: 506-455 8256
Fax: 506-459 2829

Charles Emmrys, Private Practice

115 Connaught Avenue
Moncton, NB, E1C 3P4
Tel: 506-856-3262
Fax: 506-856-2238

The Daycentre, George Dumont Hospital

185 Church Street
Moncton, NB, E1C 5A1
Tel: 506-862-4144
Fax: 506-862-4322

Eating Disorder Resource Centre

New Brunswick YWCA
35 Highfield Street
Moncton, NB, E1C 5N1
Tel: 506-855-4349
Fax: 506-855-3320

Health & Community Services

77 Vaughn Harvey Blvd.
Moncton, NB, E1C 8R3
Tel: 506-856-2401

Service de Psychologie

l'Université de Moncton
C-101, Centre étudiant
Moncton, NB, E1A 3E9
Tel: 506-858-4007
Fax: 506-858-4492

PRINCE EDWARD ISLAND

Richmond Centre
PO Box 2000 STN CENTRAL
Charlottetown, PE, C1A 7N8
Tel: 902-368-4430
Fax: 902-368-4427

HUGS, Royal Trust Tower
LL100 - 119 Kent Street
Charlottetown, PE, C1A 1N3
Tel: 902-566 4847
Fax: 902-892 4433
Contact: Cheryl Turnbull

Self-Help Clearinghouse
PO Box 785 STN CENTRAL
181 Kent Street
Charlottetown, PE, C1A 7L9
Tel: 902-628-1648

Souris Teen Group
Child and Family Division
Souris Regional Service Centre
PO Box 550
Souris, PE, C0A 2B0
Tel: 902-687-7060
Fax: 902-687-7091

HUGS
153 Spring Street
Summerside, PE, C1N 3G2
Tel: 902-436 2438
Fax: 902-436 0124

NOVA SCOTIA

HUGS

Annapolis Community Health Centre
PO Box 426
Annapolis Royal, NS, B0S 1A0
Tel: 902-532 2381
Fax: 902-532 2113

St. Martha's Regional Hospital
25 Bay Street
Antigonish, NS, B2G 2G5
Tel: 902-863-4511
Fax: 902-863-4496

HUGS

Western Kings Memorial Health Centre
PO Box 490
Berwick, NS, B0P 1E0
Tel: 902-542 6310
Fax: 902-542 6333

HUGS

115 South Street
Bridgetown, NS, B0S 1C0
Tel: 902-665-4131
Fax: 902-665-4133

Health Services Association, South Shore
90 Glen Allan Drive
Bridgewater, NS, B4V 3S6
Tel: 902-527-5228
Fax: 902-543-3120

HUGS, Sacred Heart Hospital
PO Box 129
Cheticamp, NS, B0E 1H0
Tel: 902-224 2450
Fax: 902-224 2903

Eating Disorders Action Group
7 - 106 King Street
Dartmouth, NS, B2Y 2S1
Tel: 902-469-0650
Fax: 902-469-9918

HUGS, Digby General Hospital
PO Box 820
Digby, NS, B0V 1A0
Tel: 902-245-2501
Fax: 902-245-5517

HUGS, Eskasoni Health Centre, RR#2
East Bay, NS, B0A 1H0
Tel: 902-379 2666
Fax: 902-379 2172

Eating Disorder Resource Network
55 - 5222 Green Street
Halifax, NS, B3H 1N7
Tel: 902-425-0345

Eating Disorder Clinic, Queen Elizabeth
II Health Science Centre
Third Floor, Lane Building
Camp Hill Site
Halifax, NS, B3H 2E2
Tel: 902-473-6288
Fax: 902-473-6282

Eating Disorder Program, Victoria
General Hospital
1278 Tower Road
Halifax, NS, B3H 2Y9
Tel: 902-473-6285 or 902-428-2110

Eating Disorder Resource Network, Self-
Help Eating Disorder Awareness Group
6235 St. Mathais Street
Halifax, NS, B3L 2S4
Tel: 902-423-0360

IWK-Grace Health Services Centre
Halifax, NS, B3J 3G9
Tel: 902-428-8409
Fax: 902-428-8736

Psychological and Counselling Services
Fourth Floor, Student Union Building
Dalhousie University
6136 University Avenue
Halifax, NS, B3H 4J2
Tel: 902-494-2081
Fax: 902-494-1984

Body Image and Eating Program for
Women
Twin Oaks Continuing Care Centre
RR# 2
Musquodoboit Harbour, NS, B0J 2L0
Tel: 902-889-2200
Fax: 902-889-2200

HUGS, Strait Richmond Hospital
RR# 1 Cleveland
Richmond County, NS, B0E 1J0
Tel: 902-625 3100
Fax: 902-625 3804

Eating Disorder Program, Cape Breton
Cape Breton Healthcare Complex
1482 George Street
Sydney, NS, B1P 1P3
Tel: 902-567-8000
Fax: 902-567-7905

HUGS, Colchester Regional Hospital
207 Willow Street
Truro, NS, B2N 5A1
Tel: 902-893 4321 ex 129
Fax: 902-893 5533

HUGS
PO Box 146
Windsor, NS, B0N 2T0
Tel: 902-798 2358
Fax: 902-798 4435

Mental Health Clinic
Hants Community Hospital
9 Payzant Drive
Windsor, NS, B0N 2T0
Tel: 902-792-2042
Fax: 902-798-0709

Outpatient Nutrition Services
Hants Community Hospital
9 Payzant Drive
Windsor, NS, B0N 2T0
Tel: 902-792-2000 or 902-792-2052
Fax: 902-798-4435

HUGS
E.K.M. Community Health Center
PO Box 1180
Wolfville, NS, B0P 1X0
Tel: 902-542 2266
Fax: 902-542 4619

HUGS
Site 16, Comp A9, RR#2
Wolfville, NS, B0P 8B3
Tel: 902-542 4055

Girls in the 90's, Public Health Services
60 Vancouver Street
Yarmouth, NS, B5A 2P5
Tel: 902-742-7141
Fax: 902-742-6062

Yarmouth Mental Health Centre
60 Vancouver Street
Yarmouth, NS, B5A 2P5
Tel: 902-742-4222
Fax: 902-742-2320

NEWFOUNDLAND

Health Care Corporation of St. John's
c/o Leonard A. Miller Centre
100 Forest Road
St. John's, NF, A1A 1E5
Tel: 709-737-3872
Fax: 709-737-3883

HUGS, Newfoundland
School for the Deaf
425 Topsail Road
St. John's, NF, A1E 5N7
Tel: 709-364 1234
Fax: 709-729 5848

St. Clare's Mercy Hospital
154 LeMarchant Road
St. John's, NF, A1C 5B8
Tel: 709-777-5000

APPENDIX C: INTERVIEW QUESTIONS

1. Do you think that previous or current experiences of family violence are linked to eating disorders? Can you elaborate on this?
2. In your experience, have your patients/clients revealed past or present experiences of abuse?
3. In your estimation, how prevalent is this connection?
4. Can you provide an approximate range, in percentages, of the number of clients who have experienced abuse and have manifested this in the form of eating disorders? Are they predominantly male, female, or are they both? What age groups do they represent?
5. Do you factor past or present experiences of abuse into your treatment approach to eating disorders?
6. Can you elaborate on your treatment approach?